

A tight upper bound on the (2,1)-total labeling number of outerplanar graphs

Toru Hasunuma¹, Toshimasa Ishii², Hirotaka Ono³ and Yushi Uno⁴

¹ Department of Mathematical and Natural Sciences, The University of Tokushima, Tokushima 770-8502 Japan.
hasunuma@ias.tokushima-u.ac.jp

² Department of Information and Management Science, Otaru University of Commerce, Otaru 047-8501, Japan.
ishii@res.otaru-uc.ac.jp

³ Department of Computer Science and Communication Engineering, Kyushu University, Fukuoka 812-8581, Japan.
ono@csce.kyushu-u.ac.jp

⁴ Department of Mathematics and Information Sciences, Graduate School of Science, Osaka Prefecture University, Sakai 599-8531, Japan. uno@mi.s.osakafu-u.ac.jp

Abstract. A (2, 1)-total labeling of a graph G is an assignment f from the vertex set $V(G)$ and the edge set $E(G)$ to the set $\{0, 1, \dots, k\}$ of nonnegative integers such that $|f(x) - f(y)| \geq 2$ if x is a vertex and y is an edge incident to x , and $|f(x) - f(y)| \geq 1$ if x and y are a pair of adjacent vertices or a pair of adjacent edges, for all x and y in $V(G) \cup E(G)$. The (2, 1)-total labeling number $\lambda_2^T(G)$ of a graph G is defined as the minimum k among all possible assignments. In [4], Chen and Wang conjectured that all outerplanar graphs G satisfy $\lambda_2^T(G) \leq \Delta(G) + 2$, where $\Delta(G)$ is the maximum degree of G , while they also showed that it is true for G with $\Delta(G) \geq 5$. In this paper, we solve their conjecture completely, by proving that $\lambda_2^T(G) \leq \Delta(G) + 2$ even in the case of $\Delta(G) \leq 4$.

1 Introduction

In the channel/frequency assignment problems, we need to assign different frequencies to ‘close’ transmitters so that they can avoid interference. The $L(p, q)$ -labelings of a graph have been widely studied as one of important graph theoretical models of this problem. An $L(p, q)$ -labeling of a graph G is an assignment f from the vertex set $V(G)$ to the set $\{0, 1, \dots, k\}$ of nonnegative integers such that $|f(x) - f(y)| \geq p$ if x and y are adjacent and $|f(x) - f(y)| \geq q$ if x and y are at distance 2, for all x and y in $V(G)$. We call this invariant (i.e., the minimum value k) the $L(p, q)$ -labeling number and denote it by $\lambda_{p,q}(G)$. Notice that we can use $k + 1$ different labels when $\lambda_{p,q}(G) = k$ since we can use 0 as a label for conventional reasons. We can find many results on $L(p, q)$ -labelings in comprehensive surveys by Calamoneri [3] and by Yeh [16].

In [15], Whittlesey et al. studied the $L(2, 1)$ -labeling number of incidence graphs, where the *incidence graph* of a graph G is the graph obtained from G by replacing each edge (v_i, v_j) with two edges (v_i, v_{ij}) and (v_j, v_{ij}) introducing one new vertex v_{ij} . Observe that an $L(p, 1)$ -labeling of the incidence graph of a given graph G can be regarded as an assignment f from $V(G) \cup E(G)$ to the set $\{0, 1, \dots, \ell\}$ of nonnegative integers such that $|f(x) - f(y)| \geq p$ if x is a vertex and y is an edge incident to x , and $|f(x) - f(y)| \geq 1$ if x and y are a pair of adjacent vertices or a pair of adjacent edges, for all x and y in $V(G) \cup E(G)$. Such a labeling of G is called a $(p, 1)$ -total labeling of G , introduced by Havet and Yu [7,8]. The $(p, 1)$ -total labeling number is defined as the minimum value ℓ , and denoted by $\lambda_p^T(G)$.

Notice that a $(1, 1)$ -total labeling of G is equivalent to a total coloring of G . Generalizing the Total Coloring Conjecture [2,13], Havet and Yu [7,8] conjectured that

$$\lambda_p^T(G) \leq \Delta(G) + 2p - 1 \quad (1)$$

holds for any graph G , where $\Delta(G)$ denotes the maximum degree of a vertex in G . They also showed that (i) $\lambda_p^T(G) \leq \min\{2\Delta(G) + p - 1, \chi(G) + \chi'(G) + p - 2\}$ for any graph G where $\chi(G)$ and $\chi'(G)$ denote the chromatic number and the chromatic index of G , respectively, (ii) $\lambda_2^T(G) \leq 2\Delta(G)$ if $p = 2$ and $\Delta(G) \geq 2$, (iii) $\lambda_2^T(G) \leq 2\Delta(G) - 1$ if $p = 2$ and $\Delta(G)$ is an odd ≥ 5 , and (iv) $\lambda_p^T(G) \leq n + 2p - 2$ if G is the complete graph where $n = |V(G)|$; the conjecture (1) is true if (a) $p \geq \Delta(G)$, (b) $p = 2$ and $\Delta(G) \leq 3$, or (c) G is the complete graph. By (i), it follows that $\lambda_p^T(G) \leq \Delta(G) + p$ for any bipartite graph [1,7,8] (by $\chi(G) \leq 2$ and $\chi'(G) = \Delta(G)$ [9]). Also, Bazzaro et al. [1] investigated that $\lambda_p^T(G) \leq \Delta(G) + p + s$ for any s -degenerated graph (by $\chi(G) \leq s + 1$ and $\chi'(G) \leq \Delta(G) + 1$ [12]), where an s -degenerated graph G is a graph which can be reduced to a trivial graph by successive removal of vertices with degree at most s , and that $\lambda_p^T(G) \leq \Delta(G) + p + 3$ for any planar graph (by

the Four-Color Theorem). They also showed sufficient conditions about $\Delta(G)$ and girth for which the conjecture (1) holds. In [11], Montassier and Raspaud proved that $\lambda_p^T(G) \leq \Delta(G) + 2p - 2$ when $p \geq 2$ and $\Delta(G)$ and the maximum average of G degree satisfy some conditions. In [10], Lih et al. showed that $\lambda_p^T(G) \leq \lfloor 3\Delta(G)/2 \rfloor + 4p - 3$ for any graph G . They also investigated $\lambda_p^T(K_{m,n})$ of the complete bipartite graphs $K_{m,n}$.

Let G be an outerplanar graph. In [1], Bazzaro et al. pointed out that $\lambda_p^T(G) \leq \Delta(G) + p + 1$ for any outerplanar graph other than an odd cycle, since any outerplanar graph is 2-degenerated, and any outerplanar graph other than an odd cycle satisfies $\chi'(G) = \Delta(G)$ [5]. In particular, Chen and Wang [4] conjectured that in the case of $p = 2$, $\lambda_2^T(G) \leq \Delta(G) + 2$, which is a better bound than the conjecture (1). They also proved that this conjecture is true if (i) $\Delta(G) \geq 5$, (ii) $\Delta(G) = 3$ and G is 2-connected, or (iii) $\Delta(G) = 4$ and every two faces consisting of three vertices have no vertex in common. In the case of $\Delta(G) = 2$ (i.e., G is a path or a cycle), we can easily see that $\lambda_2^T(G) \leq 4$, since the incidence graph of a path (resp., a cycle) is also a path (resp., a cycle), and the $L(2, 1)$ -labeling number $\lambda_{2,1}(C_n)$ for a cycle C_n with n vertices is at most 4 [6]. The cases of $\Delta(G) \in \{0, 1\}$ are clear. However, the general cases of $\Delta(G) \in \{3, 4\}$ were left open. In this paper, we solve Cheng and Wang's conjecture completely, by showing the remaining cases of $\Delta(G) \in \{3, 4\}$. On the other hand, the bound $\Delta(G) + 2$ is tight, since there exist infinitely many outerplanar graphs G such that $\lambda_2^T(G) = \Delta(G) + 2$ if $\Delta(G) \geq 2$, as investigated in [4].

The rest of this paper is organized as follows. Section 2 gives basic definitions. In Sections 3 and 4, we show that $\lambda_2^T(G) \leq \Delta(G) + 2$ in the cases of $\Delta(G) = 3$ and $\Delta(G) = 4$, respectively.

2 Preliminaries

A graph G is an ordered set of its vertex set $V(G)$ and edge set $E(G)$ and is denoted by $G = (V(G), E(G))$. We assume throughout this paper that a graph is undirected, simple and connected, unless otherwise stated. Therefore, an edge $e \in E(G)$ is an unordered pair of vertices u and v , which are *end vertices* of e , and we often denote it by $e = (u, v)$. For two graphs G_1 and G_2 , we denote $(V(G_1) \cup V(G_2), E(G_1) \cup E(G_2))$ by $G_1 + G_2$. For a vertex set $V' \subseteq V(G)$, let $G - V'$ be the subgraph of G induced by $V(G) - V'$. For an edge set E' , let $G - E'$ (resp., $G + E'$) be the graph $(V, E - E')$ (resp., $(V \cup V(E'), E \cup E')$ where $V(E')$ is the set of end vertices of edges in E'). Two vertices u and v are *adjacent* if $(u, v) \in E(G)$. Let $N_G(v)$ denote the set of neighbors of a vertex v in G ; $N_G(v) = \{u \in V \mid (u, v) \in E(G)\}$. The *degree* of a vertex v is $|N_G(v)|$, and is denoted by $d_G(v)$. A vertex v with $d_G(v) = k$ is called a *k-vertex*. We use $\Delta(G)$ (resp., $\delta(G)$) to denote the maximum (resp., minimum) degree of a graph G . A *path* P in G is a sequence $v_1, v_2, \dots, v_\ell$ of vertices such that $(v_i, v_{i+1}) \in E(G)$ for $i = 1, 2, \dots, \ell - 1$; we may denote P as $(v_1, v_2, \dots, v_\ell)$ and let $V(P) = \{v_1, v_2, \dots, v_\ell\}$ and $E(P) = \{(v_1, v_2), (v_2, v_3), \dots, (v_{\ell-1}, v_\ell)\}$. We often drop G in these notations if there are no confusions.

The *vertex-connectivity* of a graph G , denoted by $\kappa(G)$, is defined as the minimum cardinality of a vertex set $V' \subseteq V(G)$ such that $G - V'$ is disconnected or trivial. A vertex v is called a *cut vertex* (of G) if $G - v$ is disconnected. Notice that a graph G has no cut vertex if and only if $\kappa(G) \geq 2$. Two vertices u and u' are called *2-connected* if they belong to the same connected component in $G - v$ for any cut vertex v of G with $v \notin \{u, u'\}$. A subgraph G' of G is called a *2-connected component* if every two vertices in G' are 2-connected and G' is maximal to this property. It is not difficult to see that if $\kappa(G) = 1$, then there exists a 2-connected component which contains exactly one cut vertex of G .

A graph is called *planar* if it can be drawn in the plane without generating a crossing by two edges. For such a drawing of G in the plane, after omitting the line segments, the plane is divided into regions; such a region is called a *face*. A face whose vertex set is $\{u_1, u_2, \dots, u_k\}$ with $(u_i, u_{i+1}) \in E(G)$, $i = 1, 2, \dots, k$ (where $u_{k+1} = u_1$) is denoted by $[u_1 u_2 \dots u_k]$. We call a face consisting of k vertices a *k-face*.

A planar graph G is called *outerplanar* if it can be drawn in the plane so that all vertices lie on the boundary of some face. Such a drawing is referred to as an *outerplane graph*. The face whose boundary contains all vertices of G is called the *outer face* and all other faces are called *inner faces*. We call an edge belonging to the boundary of the outer face an *outer edge* and all other edges *inner edges*. An inner face is called an *endface* if its boundary contains exactly one inner edge (note that there exist at least two endfaces if G contains an inner edge).

Let $\mathcal{L}_k = \{0, 1, \dots, k\}$ and $f : V(G) \cup E(G) \rightarrow \mathcal{L}_k$ be a $(2, 1)$ -total labeling of G . For an edge $e = (u, v) \in E(G)$, we may denote $f(e)$ by $f(uv)$. Let $\bar{f}$ denote the labeling such that $\bar{f}(z) = k - f(z)$ for each $z \in V(G) \cup E(G)$. Notice that $\bar{f}$ is also a $(2, 1)$ -total labeling of G .

3 Case $\Delta(G) = 3$

Let $G = (V, E)$ be an outerplane graph with $\Delta(G) = 3$. Chen and Wang [4] showed that if $\kappa(G) \geq 2$, then $\lambda_2^T(G) \leq 5$. First review their constructive proof of the case of $\kappa(G) \geq 2$. Assume that on the boundary of its

outerface, each 3-vertex (i.e., each vertex which some inner edge is incident to) is numbered $x_1, x_2, \dots, x_p$ in clockwise order. Also assume that for $i = 1, 2, \dots, p$, each 2-vertex on the path connecting x_i and x_{i+1} (on the boundary of its outerface) is numbered $y_1^i, y_2^i, \dots, y_{q_i}^i$ in clockwise order, where $x_{p+1} = x_1$. A (2,1)-total labeling $f : V \cup E \rightarrow \mathcal{L}_5$ can be obtained by the following algorithm, which slightly generalizes Chen and Wang's method.

Algorithm LABEL-K2

Input: An outerplane graph G with $\kappa(G) \geq 2$.

Output: A total (2,1)-labeling $f : V \cup E \rightarrow \mathcal{L}_5$ of G .

1. Assign labels 0 and 1 alternately to a sequence $x_1, x_2, \dots, x_p$ of 3-vertices.
2. For each $i = 1, 2, \dots, p$, label 2-vertices as follows. Without loss of generality, let $f(x_i) = 0$ and $f(x_{i+1}) = 1$.
 - 2.1. If q_i is even, then assign labels 1 and 0 alternately to a sequence $y_1^i, y_2^i, \dots, y_{q_i}^i$ of 2-vertices.
 - 2.2. If q_i is odd, then assign label 2 to y_j^i for some j , and labels 1 and 0 alternately to a sequence $y_1^i, y_2^i, \dots, y_{q_i}^i$ of 2-vertices other than y_j^i .
3. Assign label 3 to each inner edge.
4. Label outer edges as follows.
 - 4.1. If $|V(G)|$ is even, then assign labels 4 and 5 alternately in clockwise order.
 - 4.2. If $|V(G)|$ is odd, then pick up an arbitrary endface $F = [x_i y_1^i y_2^i \dots y_{q_i}^i x_{i+1}]$. Without loss of generality, let $f(x_i) = 0$ and $f(x_{i+1}) = 1$.
 - 4.2.1. If q_i is an even or an odd with $q_i \geq 3$, then assign label 3 to (y_1^i, y_2^i) and labels 4 and 5 alternately to all remaining outer edges. Then if $f(y_1^i) = 2$ or $f(y_2^i) = 2$, then reassign labels for y_1^i, y_2^i , and y_3^i as $f(y_1^i) := 1, f(y_2^i) := 0$, and $f(y_3^i) := 2$.
 - 4.2.2. If $q_i = 1$, then assign labels 5, 2, and 3 to $y_1^i, (x_i, y_1^i)$, and (y_1^i, x_{i+1}) , respectively, and labels 4 and 5 alternately to all remaining outer edges. Then if the label for two outer edges incident to x_i and x_{i+1} other than (x_i, y_1^i) and (y_1^i, x_{i+1}) is 4 (resp., 5), reassign a label for (x_i, x_{i+1}) as $f(x_i x_{i+1}) := 5$ (resp., 4).

Observe that for each inner edge (x_i, x_j) , we have $f(x_i) \neq f(x_j)$, since otherwise the subgraph G' of G induced by $V(P_i) \cup V(P_{i+1}) \cup \dots \cup V(P_{j-1})$ would have odd 3-vertices, a contradiction, where P_i denotes the path $(x_i, y_1^i, \dots, y_{q_i}^i, x_{i+1})$. Notice that the reassignment in Step 4.2.1 is necessary because otherwise $|f(y_1^i) - f(y_2^i)| \leq 1$ or $|f(y_2^i) - f(y_3^i)| \leq 1$ would hold. Also notice that in Step 4.2.2, $f(e) = f(e')$ holds for two outer edges e (resp., e') incident to x_i (resp., x_{i+1}) other than (x_i, y_1^i) (resp., (y_1^i, x_{i+1})), since $|V(G)|$ is odd and $q_i = 1$. Thus, a labeling f obtained from the above procedure is a (2,1)-total labeling of G . Here, we have the following easy observation:

Observation 1. In Step 4.2, we can start with assigning an arbitrary label in $\{4, 5\}$ to an arbitrary outer edge for the label assignment of outer edges except (y_1^i, y_2^i) (resp., (x_i, y_1^i) and (y_1^i, x_{i+1})) in Step 4.2.1 (resp., Step 4.2.2).

Moreover, we can see the following property.

Lemma 1. Let $G = (V, E)$ be an outerplane graph with $\Delta(G) = 3$, G_1 be a 2-connected component in G , and (u, v) be an inner edge in G_1 . Assume that $G_1 - \{u, v\}$ contains a connected component G_2 with no cut vertex in G and $|V(G_2)| \geq 2$. Let $u' \in V(G_2)$ (resp., $v' \in V(G_2)$) be the neighbor of u (resp., v) in G (note that by $\Delta(G) = 3$, such u' and v' are uniquely determined and that by $|V(G_2)| \geq 2$, we have $u' \neq v'$). If $G' \triangleq (G - V(G_2)) + \{(u, u'), (v, v')\}$ has a (2,1)-total labeling $f : V(G') \cup E(G') \rightarrow \mathcal{L}_5$ such that

- (i) $\{f(u'), f(v')\} \subseteq \{0, 1\}$ and $\{f(uu'), f(vv')\} \subseteq \{4, 5\}$ or
- (ii) $\{f(u'), f(v')\} \subseteq \{4, 5\}$ and $\{f(uu'), f(vv')\} \subseteq \{0, 1\}$,

then there exists a (2,1)-total labeling $g : V(G) \cup E(G) \rightarrow \mathcal{L}_5$.

Proof. By symmetry of labelings, we only consider the case where $f(u') \neq f(v')$, $\{f(u'), f(v')\} \subseteq \{0, 1\}$, and $\{f(uu'), f(vv')\} \subseteq \{4, 5\}$. Without loss of generality, let $f(u') = 0$ and $f(v') = 1$. There are the two possible cases (Case-1) $f(uu') = f(vv')$ and (Case-2) $f(uu') \neq f(vv')$.

(Case-1) Let G_3 be the graph obtained from G_2 by adding two new vertices x and y and three new edges (u', x) , (x, y) , and (y, v') . Notice that G_3 is an outerplane graph with $\kappa(G_3) \geq 2$. There are the following two possible cases: (1.1) both of u' and v' are 2-vertices or 3-vertices, (1.2) otherwise.

(1.1) If u' and v' are both 2-vertices, then add an inner edge (u', v') to G_3 and redefine G_3 by the resulting graph. By applying Algorithm LABEL-K2 to G_3 as $x_1 := u'$, we can obtain a (2,1)-total labeling $f_1 : V(G_3) \cup$

$E(G_3) \rightarrow \mathcal{L}_5$ such that $f_1(u') = 0$ and $f_1(v') = 1$. Moreover, by choosing an endface not containing $\{x, y\}$ in Step 4.2 of Algorithm LABEL-K2, we can obtain such a labeling f_1 that $f_1(xu') = f_1(yv') = f(uu') (= f(vv'))$ (note that by Observation 1, we can start with assigning label $f(uu')$ to (x, u') for the label assignment of outer edges). Observe that the labeling g such that $g(z) = f(z)$ for each $z \in V(G') \cup E(G')$ and $g(z) = f_1(z)$ for each $z \in V \cup E - (V(G') \cup E(G'))$ is a (2,1)-total labeling of G .

(1.2) Assume that u' is a 3-vertex and v' is a 2-vertex (the other case can be treated similarly). Let $f_1 : V(G_3) \cup E(G_3) \rightarrow \mathcal{L}_5$ be a (2,1)-total labeling obtained by applying Algorithm LABEL-K2 to G_3 as $x_1 := u'$; $f_1(u') = 0$. Moreover, as observed in (Case-1.1), we can obtain such a f_1 that $f_1(xu') = f_1(yv') = f(uu')$. Also, if v' is not adjacent to a 3-vertex, we can obtain such a f_1 that $f_1(v') = 1$ by avoiding assigning label 2 to any vertex in $\{x, y, v'\}$ in Step 2.2. Observe that if $f_1(v') = 1$, then a (2,1)-total labeling g of G can be constructed by combining f and f_1 similarly to (Case-1.1). Hence, we consider the case where the vertex $w (\neq y)$ adjacent to v' in G_3 is a 3-vertex; $f_1(w) = 1$. Then we can reassign labels for x, y , and v' as any one of two possible labelings (a) $f_1(x) = 1, f_1(y) = 0$, and $f_1(v') = 2$ and (b) $f_1(x) = 1, f_1(y) = 2$, and $f_1(v') = 0$ without violating the feasibility. Observe that the labeling g such that $g(z) = f(z)$ for each $z \in V(G') \cup E(G') - \{v'\}$, $g(z) = f_1(z)$ for each $z \in V \cup E - (V(G') \cup E(G'))$, and $g(v') = \{0, 2\} - \{f(v)\}$ is a (2,1)-total labeling of G .

(Case-2) Let G_3 be the graph obtained from G_2 by adding a new vertex x and two new edges (u', x) and (x, v') . Notice that G_3 is an outerplane graph with $\kappa(G_3) \geq 2$. There are the following two possible cases: (2.1) both of u' and v' are 2-vertices or 3-vertices, (2.2) otherwise.

(2.1) If u' and v' are both 2-vertices, then add an inner edge (u', v') to G_3 and redefine G_3 by the resulting graph. By applying Algorithm LABEL-K2 to G_3 as $x_1 := u'$, we can obtain a (2,1)-total labeling $f_1 : V(G_3) \cup E(G_3) \rightarrow \mathcal{L}_5$ such that $f_1(u') = 0$ and $f_1(v') = 1$. Moreover, by choosing an endface not containing x in Step 4.2 of Algorithm LABEL-K2, we can obtain such a labeling f_1 that $f_1(xu') = f(uu')$ and $f_1(xv') = f(vv')$. Observe that the labeling g such that $g(z) = f(z)$ for each $z \in V(G') \cup E(G')$ and $g(z) = f_1(z)$ for each $z \in V \cup E - (V(G') \cup E(G'))$ is a (2,1)-total labeling of G .

(2.2) Assume that u' is a 3-vertex and v' is a 2-vertex (the other case can be treated similarly). Let $f_1 : V(G_3) \cup E(G_3) \rightarrow \mathcal{L}_5$ be a (2,1)-total labeling obtained by applying Algorithm LABEL-K2 to G_3 as $x_1 := u'$; $f_1(u') = 0$. Moreover, as observed in (Case-2.1), we can obtain such a f_1 that $f_1(xu') = f(uu')$ and $f_1(xv') = f(vv')$. Also, in the case where Step 2.2 is applied to the sequence of 2-vertices containing x and v' , we can assign label 1 to v' by assigning label 2 to x (notice that in this case, v' is not adjacent to any 3-vertex because the sequence of 2-vertices containing v' consists of odd vertices). Observe that if $f_1(v') = 1$, then a (2,1)-total labeling g of G can be constructed by combining f and f_1 similarly to (Case-2.1). Hence, we consider the case where Step 2.1 is applied to the sequence of 2-vertices containing x and v' ; $f_1(v') = 0$. Then we can reassign label 2 to v' without violating the feasibility. We can see that the labeling g such that $g(z) = f(z)$ for each $z \in V(G') \cup E(G') - \{v'\}$, $g(z) = f_1(z)$ for each $z \in V \cup E - (V(G') \cup E(G'))$, and $g(v') = \{0, 2\} - \{f(v)\}$ is a (2,1)-total labeling of G . $\square$

By using this lemma, we can prove that Chen and Wang's conjecture is true in the case of $\Delta(G) = 3$.

Theorem 1. *If $G = (V, E)$ is an outerplane graph with $\Delta(G) = 3$, then $\lambda_2^T(G) \leq 5$.*

Proof. We prove this by induction on $k = |V(G)| + |E(G)|$. The theorem clearly holds if $k \leq 7$. Consider the case of $k \geq 8$ and assume that for each $k' < k$, this theorem holds. We also assume that G is connected, since otherwise we can treat each component separately. Thus, $1 \leq \delta(G) \leq 2$.

Consider the case where $\delta(G) = 1$. Let u_1 be a vertex with $d(u_1) = 1$. By the induction hypothesis, $G - u_1$ has a (2,1)-total labeling $f : V(G - u_1) \cup E(G - u_1) \rightarrow \mathcal{L}_5$. Let u_2 be the neighbor of u_1 in G and u_3, u_4 be vertices adjacent to u_2 in $G - u_1$ where $u_3 = u_4$ may occur (note that $\Delta(G) = 3$). Hence we can extend f to the edge (u_1, u_2) and the vertex u_1 as follows: assign a label $a \in \mathcal{L}_5 - \{f(u_2) - 1, f(u_2), f(u_2) + 1, f(u_2u_3), f(u_2u_4)\}$ to (u_1, v_1) , and then a label in $\mathcal{L}_5 - \{f(u_2), a - 1, a, a + 1\}$ to u_1 .

Consider the case of $\delta(G) = 2$. In [4], Chen and Wang showed that if $\kappa(G) \geq 2$, then $\lambda_2^T(G) \leq 5$. Hence, we here only consider the case of $\kappa(G) = 1$. Let G_1 be a 2-connected component in G which has exactly one cut vertex v_c of G . By $\delta(G) = 2$, we have $|V(G_1)| \geq 3$ and hence $d_{G_1}(v_c) \geq 2$. By $\kappa(G) = 1$ and $\Delta(G) = 3$, it follows that $d_{G_1}(v_c) = 2$ holds and $V(G_1)$ and $V - V(G_1)$ are connected by a bridge incident to v_c , where a *bridge* is an edge whose deletion makes the graph disconnected; denote this bridge by (v_c, w) where $w \in V - V(G_1)$. By the induction hypothesis, $H = (G - V(G_1)) \cup \{(v_c, w)\}$ has a (2,1)-total labeling $f : V(H) \cup E(H) \rightarrow \mathcal{L}_5$. By symmetry, it suffices to consider the following three possible cases: (Case-1) $f(v_cw) = 5$, (Case-2) $f(v_cw) = 4$, and (Case-3) $f(v_cw) = 3$. In each case, we will extend f to a (2,1)-total labeling $f' : V \cup E \rightarrow \mathcal{L}_5$ of G . Let $f'(x) := f(x)$ for every $x \in V(H) \cup E(H)$.

First consider the case where G_1 has no inner edge. Assume that on the boundary of its outerface in G_1 , each vertex is numbered $u_1 (= v_c), u_2, \dots, u_r$ in clockwise order. Then, in each case of (Case-1)–(Case-3), we can obtain a (2,1)-total labeling f' of G in the following manner:

1. If $f'(v_c w) = 3$, $r = 3$, and $f'(w) \neq 0$, then let $f'(v_c) = 0$, $f'(u_2) := 1$, $f'(u_3) := 5$, $f'(v_c u_2) := 5$, $f'(u_2 u_3) := 3$, and $f'(u_3 v_c) := 2$. If $f'(v_c w) = 3$, $r = 3$, and $f'(w) = 0$, then let $f'(v_c) = 5$, $f'(u_2) := 2$, $f'(u_3) := 0$, $f'(v_c u_2) := 0$, $f'(u_2 u_3) := 5$, and $f'(u_3 v_c) := 2$.
2. Unless $f'(v_c w) = 3$ and $r = 3$, then assign labels for vertices or edges in G_1 as follows.
 - 2.1. Assign a label in $\{0, 1\} - \{f'(w)\}$ (say, 0) to v_c .
 - 2.2. Assign labels 1 and 0 alternately to $u_2, u_3, \dots, u_r$. Then, if r is odd, then reassign label 2 to u_r .
 - 2.3. If $f'(v_c w) \in \{3, 4\}$ (resp., $f'(v_c w) = 5$), then assign label 5 (resp., 4) to (u_r, v_c) , and labels 4 and 5 (resp., 5 and 4) alternately to the sequence $(u_r, u_{r-1}), (u_{r-1}, u_{r-2}), \dots, (u_2, v_c)$ of outer edges in counter-clockwise order. Then if $f'(v_c w) \in \{4, 5\}$, then reassign label 3 to (u_2, v_c) , and if $f'(v_c w) = 3$ and r is odd, then reassign label 3 to (u_3, u_2) and label 4 to (u_2, v_c) .

Next consider the case where G_1 has an inner edge. Assume that on the boundary of its outerface in G_1 , each 3-vertex is numbered $x_1, x_2, \dots, x_p$ in clockwise order. Also assume that for $i = 1, 2, \dots, p$, each 2-vertex on the path connecting x_i and x_{i+1} (on the boundary of its outerface) is numbered $y_1^i, y_2^i, \dots, y_{q_i}^i$ in clockwise order, where $x_{p+1} = x_1$. Also, if $q_i = 0$, then let $y_{q_i}^i := x_i$. Let P_i denote the path $(x_i, y_1^i, \dots, y_{q_i}^i, x_{i+1})$. Without loss of generality, assume that v_c is a 2-vertex between x_1 and x_2 in G_1 ; $v_c = y_\ell^1$ for some ℓ . Let $f_1 : V(G_1) \cup E(G_1) \rightarrow \mathcal{L}_5$ be a (2,1)-total labeling obtained by applying Algorithm LABEL-K2 to G_1 , while (A) choosing y_j^1 as a vertex other than v_c in Step 2.2 if $q_1 \geq 3$, and (B) choosing y_j^1 as a vertex not in $\{v_c\} \cup N_G(v_c)$ in Step 2.2 if (i) $q_1 \geq 5$ or (ii) $q_1 = 3$ and $v_c \neq y_1^1$, and (C) choosing an endface not containing v_c in Step 4.2.

Consider the case where v_c is adjacent to a 2-vertex in G_1 ; $q_1 > 1$. Without loss of generality, assume that $y_{\ell-1}^1$ is a 2-vertex. Then by $q_1 > 1$ and the choice of y_j^1 in Step 2.2 of Algorithm LABEL-K2, we can see that $f_1(v_c) \neq 2$; $f_1(v_c) \in \{0, 1\}$. Moreover, we can assume that $f_1(v_c) \in \{0, 1\} - \{f'(w)\}$ since otherwise we recompute a (2,1)-total labeling f'_1 of G_1 by starting with x_2 instead of x_1 in Step 1 of Algorithm LABEL-K2 and redefine f_1 by f'_1 . Also, we can assume that in the case of $f'(v_c w) = 4$ (resp., 5), if $q_1 \neq 3$ or $v_c \neq y_1^1$, then $f_1(y_{\ell-1}^1 v_c) = 4$ (resp., 5) and if $q_1 = 3$ and $v_c = y_1^1$, then $f_1(v_c y') = 4$ (resp., 5) for $y' \in \{y_1^1, y_3^1\}$ with $f_1(y') \neq 2$ (note that this is possible by Observation 1 and that in the case of $q_1 = 3$ and $v_c = y_1^1$, exactly one of y_1^1 and y_3^1 has label 2 in f_1). Let $f'(z) := f_1(z)$ for each $z \in V(G_1) \cup E(G_1)$. Observe that if $f'(v_c w) = 3$, then f' is a (2,1)-total labeling of G . In the case of $f'(v_c w) \in \{4, 5\}$, if $q_1 \neq 3$ or $v_c \neq y_1^1$ (resp., $q_1 = 3$ and $v_c = y_1^1$), reassign a label for $(y_{\ell-1}^1, v_c)$ (resp., (y', v_c)) as $f'(y_{\ell-1}^1 v_c) := 3$ (resp., $f'(y' v_c) := 3$). Then, we can see that the resulting f' is a (2,1)-total labeling of G since if $q_1 \neq 3$ or $v_c \neq y_1^1$, then $f'(y_{\ell-1}^1) \neq 2$ by the choice of y_j^1 in Step 2.2 and otherwise then $f'(y') \neq 2$.

Finally, consider the case where $N_{G_1}(v_c) = \{x_1, x_2\}$; $q_1 = 1$. Let $x_{p'}$ be a 3-vertex such that $(x_1, x_{p'})$ is an inner edge of G_1 . Let G_2 be the subgraph of G_1 induced by $V(P_1) \cup V(P_2) \cup \dots \cup V(P_{p'-1})$. Let $u' \in V(G_1) - V(G_2)$ (resp., $v' \in V(G_1) - V(G_2)$) be the neighbor of x_1 (resp., $x_{p'}$). Notice that the component in $G - \{x_1, x_{p'}\}$ containing $\{u', v'\}$ has no cut vertex in G . Below, we will show that if $u' \neq v'$, then there exists a (2,1)-total labeling g' of $(H + G_2) + \{(x_1, u'), (x_{p'}, v')\}$ satisfying the conditions of Lemma 1. Note that in this case, Lemma 1 ensures the existence of a (2,1)-total labeling $g : V \cup E \rightarrow \mathcal{L}_5$ of G . Also, in the case of $u' = v'$, we will show directly the existence of a (2,1)-total labeling $g : V \cup E \rightarrow \mathcal{L}_5$ of G .

(Case-1) $f(v_c w) = 5$.

(1.1) Assume that $f(w) \neq 3$.

Let $f'(z) := \overline{f_1}(z)$ for each $z \in V(G_1) \cup E(G_1)$. Then, f' is a (2,1)-total labeling of G (note that $f'(v_c) = 3$).

(1.2) Assume that $f(w) = 3$.

If $p' = 2$ and $u' = v'$, then let $f'(x_1) := 5$, $f'(v_c) := 0$, $f'(x_2) := 2$, $f'(u') := 0$, $f'(x_1 v_c) := 3$, $f'(v_c x_2) := 4$, $f'(x_1 x_2) := 0$, $f'(x_2 u') := 5$, and $f'(u' x_1) := 2$. Observe that the resulting labeling f' is a (2,1)-total labeling of G .

If $p' = 2$ and $u' \neq v'$, then let $f'(x_1) := 5$, $f'(v_c) := 0$, $f'(x_2) := 4$, $f'(u') := 4$, $f'(v') := 5$, $f'(x_1 v_c) := 3$, $f'(v_c x_2) := 2$, $f'(x_1 x_2) := 0$, $f'(x_2 v') := 1$, and $f'(u' x_1) := 1$. Observe that the resulting labeling f' is feasible for $H + G_2 + \{(x_1, u'), (x_2, v')\}$ and satisfies the conditions of Lemma 1.

Assume that $p' \neq 2$. Then we can obtain a (2,1)-total labeling of G in the following manner, where “assigning a label k to $z \in V \cup E$ ” means that $f'(z) := k$.

1. Assign label 3 to each inner edge in G_2 .
2. Let $f'(v_c) := 0$. Assign labels 1 and 0 alternately to 3-vertices $x_2, x_3, \dots, x_{p'-1}$ (note that as a result, $f'(x_{p'-1}) = 0$). Assign labels to 2-vertices y_j^i ($2 \leq i \leq p' - 2$) according to Step 2 of algorithm LABEL-K2.

3. Trace outer edges from (v_c, x_2) to $(y_{q_{p'-2}}^{p'-2}, x_{p'-1})$ on $\{(v_c, x_2)\} \cup E(P_2) \cup \dots \cup E(P_{p'-2})$ in clockwise order and assign labels 4 and 5 alternately.
 - 3.1. If $q_{p'-1} > 0$, then assign labels to the remaining vertices or edges as follows. Let $f'(x_1) := 5$, $f'(x_{p'}) := 4$, $f'(x_1 v_c) := 3$, $f'(x_1 x_{p'}) := 2$, and $f'(x_{p'-1}, y_1^{p'-1}) := 2$. Assign labels 0 and 1 alternately to outer edges $(y_1^{p'-1}, y_2^{p'-1})$, $(y_2^{p'-1}, y_3^{p'-1})$, $\dots$, $(y_{q_{p'-1}}^{p'-1}, x_{p'})$, $(x_{p'}, v')$. Moreover, if $q_{p'-1}$ is even (resp., odd), then assign labels 4 and 5 (resp., 5 and 4) alternately to 2-vertices $y_1^{p'-1}, y_2^{p'-1}, \dots, y_{q_{p'-1}}^{p'-1}$. If $u' = v'$, then let $f'(u') := 3$ and $f'(u' x_1) := \{0, 1\} - \{f'(x_{p'} u')\}$. If $u' \neq v'$, then let $f'(u') := 4$, $f'(v') := 5$, and $f'(u' x_1) := 1$, and extend f' to a (2,1)-total labeling of G according to Lemma 1.
 - 3.2. If $q_{p'-1} = 0$ and $u' \neq v'$, then let $f'(x_1) := f'(v') := 4$, $f'(x_{p'}) := f'(u') := 5$, $f'(x_1 x_{p'}) := 0$, $f(x_{p'-1} x_{p'}) := f(x_1 v_c) := 2$, and $f'(x_1 u') := f'(x_{p'} v') := 1$, and extend f' to a (2,1)-total labeling of G according to Lemma 1.
 - 3.3. If $q_{p'-1} = 0$, $u' = v'$ and $f'(y_{q_{p'-2}}^{p'-2} x_{p'-1}) = 4$, then let $f'(x_1) := 5$, $f'(x_{p'}) := 3$, $f'(u') := 4$, $f'(x_1 x_{p'}) := 1$, $f(x_{p'-1} x_{p'}) := 5$, $f(x_1 v_c) := 3$, and $f'(x_1 u') := 2$, and $f'(x_{p'} u') := 0$.
 - 3.4. If $q_{p'-1} = 0$, $u' = v'$ and $f'(y_{q_{p'-2}}^{p'-2} x_{p'-1}) = 5$, then let $f'(x_1) := 5$, $f'(x_{p'}) := 2$, $f'(u') := 0$, $f'(x_1 x_{p'}) := 0$, $f(x_{p'-1} x_{p'}) := 4$, $f(x_1 v_c) := 3$, and $f'(x_1 u') := 2$, and $f'(x_{p'} u') := 5$.

(Case-2) $f(v_c w) = 4$.

(2.1) Assume that $f(w) \neq 0$.

If $p' = 2$ and $u' = v'$, then let $f'(x_1) := 5$, $f'(v_c) := 0$, $f'(x_2) := 2$, $f'(u') = 0$, $f'(x_1 v_c) := 3$, $f'(v_c x_2) := 5$, $f'(x_1 x_2) := 0$, $f'(x_2 u') := 4$, and $f'(u' x_1) := 2$.

If $p' = 2$ and $u' \neq v'$, then let $f'(x_1) := f'(v') := 5$, $f'(v_c) := 0$, $f'(x_2) := f'(u') := 4$, $f'(x_1 v_c) := 3$, $f'(v_c x_2) := 2$, $f'(x_1 x_2) := 0$, $f'(x_2 v') := 1$, $f'(u' x_1) := 1$. Observe that the resulting labeling f' is feasible for $H + G_2 + \{(x_1, u'), (x_2, v')\}$ and satisfies the conditions of Lemma 1.

The case of $p' \neq 2$ can be treated similarly to (Case-1.2), by starting with assigning to (v_c, x_2) label 5 instead of label 4.

(2.2) Assume that $f(w) = 0$.

If $p' = 2$ and $u' = v'$, then let $f'(v_c) := 1$, $f'(x_1) := 5$, $f'(x_2) := 3$, $f'(u') := 4$, $f'(x_1 v_c) := 3$, $f'(v_c x_2) := 5$, $f'(x_1 x_2) := 0$, $f'(x_1 u') := 2$, and $f'(x_2 u') := 1$.

If $p' = 2$ and $u' \neq v'$, then $f'(v_c) := 1$, $f'(x_1) := 5$, $f'(x_2) := 3$, $f'(u') := 4$, $f'(v') := 5$, $f'(x_1 v_c) := 3$, $f'(v_c x_2) := 5$, $f'(x_1 x_2) := 0$, $f'(x_1 u') := 1$, and $f'(x_2 v') := 1$. Observe that the resulting labeling is feasible for $H + G_2 + \{(x_1, u'), (x_2, v')\}$ and satisfies the conditions of Lemma 1.

Assume that $p' \neq 2$. Then we can obtain a (2,1)-total labeling of G in the following manner.

1. Assign label 3 to each inner edge in G_2 .
2. Trace outer edges from (v_c, x_2) to $(y_{q_{p'-1}}^{p'-1}, x_{p'})$ on $\{(v_c, x_2)\} \cup E(P_2) \cup \dots \cup E(P_{p'-1})$ in clockwise order and assign labels 5 and 4 alternately.
 - 2.1. If $f'(y_{q_{p'-1}}^{p'-1} x_{p'}) = 4$ and $u' \neq v'$, then assign labels for the remaining vertices and edges as follows. Let $f'(v_c) := 2$. Assign labels 1 and 0 alternately to 3-vertices $x_2, x_3, \dots, x_{p'-1}$ (notice that $f'(x_{p'-1}) = 0$). Assign labels to 2-vertices y_j^i ($2 \leq i \leq p' - 2$) according to Step 2 of algorithm LABEL-K2. Let $f'(x_1) := 3$, $f'(x_{p'}) := 2$, $f'(u') := 4$, $f'(v') := 5$, $f'(x_1 v_c) := 0$, $f'(x_1 x_{p'}) := 5$, $f'(x_1 u') := 1$, and $f'(x_{p'} v') := 0$. Assign labels 1 and 0 alternately to 2-vertices $y_1^{p'-1}, y_2^{p'-1}, \dots, y_{q_{p'-1}}^{p'-1}$. Then, extend f' to a (2,1)-total labeling of G according to Lemma 1.
 - 2.2. If $f'(y_{q_{p'-1}}^{p'-1} x_{p'}) = 5$ and $u' \neq v'$, then assign labels for the remaining vertices and edges as follows. Let $f'(v_c) := 1$. Assign labels 0 and 1 alternately to 3-vertices $x_2, x_3, \dots, x_{p'-1}$ (notice that $f'(x_{p'-1}) = 1$). Assign labels to 2-vertices y_j^i ($2 \leq i \leq p' - 2$) according to Step 2 of algorithm LABEL-K2. Let $f'(x_1) := 5$, $f'(x_{p'}) := 3$, $f'(u') := 4$, $f'(v') := 5$, $f'(x_1 v_c) := 3$, $f'(x_1 x_{p'}) := 0$, $f'(x_1 u') := 1$, and $f'(x_{p'} v') := 1$. Assign labels 0 and 1 alternately to 2-vertices $y_1^{p'-1}, y_2^{p'-1}, \dots, y_{q_{p'-1}}^{p'-1}$. Then, extend f' to a (2,1)-total labeling of G according to Lemma 1.
 - 2.3. If $f'(y_{q_{p'-1}}^{p'-1} x_{p'}) = 4$ and $u' = v'$, then assign labels for the remaining vertices and edges as follows. Let $f'(v_c) := 2$. Assign labels 1 and 0 alternately to 3-vertices $x_2, x_3, \dots, x_{p'-1}$ (notice that $f'(x_{p'-1}) = 0$). Assign labels to 2-vertices y_j^i ($2 \leq i \leq p' - 2$) according to Step 2 of algorithm LABEL-K2. Let $f'(x_1) := 3$, $f'(x_{p'}) := 2$, $f'(u') := 5$, $f'(x_1 v_c) := 0$, $f'(x_1 x_{p'}) := 5$, $f'(x_1 u') := 1$, and $f'(x_{p'} u') := 0$. Assign labels 1 and 0 alternately to 2-vertices $y_1^{p'-1}, y_2^{p'-1}, \dots, y_{q_{p'-1}}^{p'-1}$.
 - 2.4. If $f'(y_{q_{p'-1}}^{p'-1} x_{p'}) = 5$ and $u' = v'$, then assign labels for the remaining vertices and edges as follows. Let $f'(v_c) := 1$. Assign labels 0 and 1 alternately to 3-vertices $x_2, x_3, \dots, x_{p'-1}$ (notice that $f'(x_{p'-1}) = 1$). Assign labels to 2-vertices y_j^i ($2 \leq i \leq p' - 2$) according to Step 2 of algorithm LABEL-K2. Let

$f'(x_1) := 5, f'(x_{p'}) := 3, f'(u') := 4, f'(x_1v_c) := 3, f'(x_1x_{p'}) := 0, f'(x_1u') := 2$, and $f'(x_{p'}u') := 1$.
Assign labels 0 and 1 alternately to 2-vertices $y_1^{p'-1}, y_2^{p'-1}, \dots, y_{q_{p'-1}}^{p'-1}$.

(Case-3) $f(v_cw) = 3$.

First consider the case where $p' = 2$ and $u' = v'$. If $f(w) \neq 5$, then let $f'(x_1) := 0, f'(v_c) := 5, f'(x_2) := 2, f'(u') := 1, f'(x_1v_c) := 2, f'(v_cx_2) := 0, f'(x_1x_2) := 5, f'(x_2u') := 4$, and $f'(u'x_1) := 3$. If $f(w) = 5$, then let $f'(x_1) := 4, f'(v_c) := 0, f'(x_2) := 2, f'(u') := 3, f'(x_1v_c) := 2, f'(v_cx_2) := 4, f'(x_1x_2) := 0, f'(x_2u') := 5$, and $f'(u'x_1) := 1$.

Next consider the case where $p' = 2$ and $u' \neq v'$. If $f(w) \neq 5$, then let $f'(x_1) := 0, f'(v_c) := 5, f'(x_2) := 2, f'(u') := 1, f'(v') := 0, f'(x_1v_c) := 2, f'(v_cx_2) := 0, f'(x_1x_2) := 5, f'(x_2v') := 4$, and $f'(u'x_1) := 4$. If $f(w) = 5$, then let $f'(x_1) := 3, f'(v_c) := 0, f'(x_2) := 5, f'(u') := 5, f'(v') := 4, f'(x_1v_c) := 5, f'(v_cx_2) := 2, f'(x_1x_2) := 0, f'(x_2v') := 1$, and $f'(u'x_1) := 1$. Observe that in both cases, the resulting labeling is feasible for $H + G_2 + \{(x_1, u'), (x_{p'}, v')\}$ and satisfies the conditions of Lemma 1.

Finally consider the case where $p' \neq 2$. We divide this case into the following two subcases: (3.1) $f'(w) \neq 0$ and (3.2) $f'(w) = 0$.

(3.1) Assume that $f'(w) \neq 0$.

We can obtain a (2,1)-total labeling of G in the following manner.

1. Assign label 3 to each inner edge in G_2 .
2. Let $f'(v_c) := 0$. Assign labels 1 and 0 alternately to 3-vertices $x_2, x_3, \dots, x_{p'-1}$ (note that as a result, $f'(x_{p'-1}) = 0$). Assign labels to 2-vertices y_j^i ($2 \leq i \leq p' - 2$) according to Step 2 of algorithm LABEL-K2. Assign labels 1 and 0 alternately to 2-vertices $y_1^{p'-1}, y_2^{p'-1}, \dots, y_{q_{p'-1}}^{p'-1}$.
3. Trace outer edges from (v_c, x_2) to $(y_{q_{p'-1}}^{p'-1}, x_{p'})$ on $\{(v_c, x_2)\} \cup E(P_2) \cup \dots \cup E(P_{p'-1})$ in clockwise order and assign labels 4 and 5 (resp., 5 and 4) alternately if the number of outer edges in G_2 is even (resp., odd). Notice that as a result, the edge $(y_{q_{p'-1}}^{p'-1}, x_{p'})$ has label 5.
 - 3.1. If $u' \neq v'$, then let $f'(x_1) := 5, f'(x_{p'}) := 3, f'(u') := 4, f'(v') := 5, f'(x_1v_c) := 2, f'(x_1x_{p'}) := 0, f'(x_1u') := 1$, and $f'(x_{p'}v') := 1$. Then, extend f' to a (2,1)-total labeling of G according to Lemma 1.
 - 3.2. If $u' = v'$, then let $f'(x_1) := 5, f'(x_{p'}) := 2, f'(u') := 0, f'(x_1v_c) := 2, f'(x_1x_{p'}) := 0, f'(x_1u') := 3$, and $f'(x_{p'}u') := 4$.

(3.2) Assume that $f'(w) = 0$.

We can obtain a (2,1)-total labeling of G in the following manner.

1. Assign label 2 to each inner edge in G_2 .
2. Let $f'(v_c) := 5$. Assign labels 4 and 5 alternately to 3-vertices $x_2, x_3, \dots, x_{p'-1}$ (note that as a result, $f'(x_{p'-1}) = 5$). Assign labels to 2-vertices y_j^i ($2 \leq i \leq p' - 2$) according to Step 2 of Algorithm LABEL-K2', where Algorithm LABEL-K2' denotes the algorithm obtained from Algorithm LABEL-K2 by replacing each label i to be assigned with $5 - i$. Assign labels 4 and 5 alternately to 2-vertices $y_1^{p'-1}, y_2^{p'-1}, \dots, y_{q_{p'-1}}^{p'-1}$.
3. Trace outer edges from (v_c, x_2) to $(y_{q_{p'-1}}^{p'-1}, x_{p'})$ on $\{(v_c, x_2)\} \cup E(P_2) \cup \dots \cup E(P_{p'-1})$ in clockwise order and assign labels 1 and 0 (resp., 0 and 1) alternately if the number of outer edges in G_2 is even (resp., odd). Notice that as a result, the edge $(y_{q_{p'-1}}^{p'-1}, x_{p'})$ has label 0.
 - 3.1. If $u' \neq v'$, then let $f'(x_1) := 0, f'(x_{p'}) := 2, f'(u') := 1, f'(v') := 0, f'(x_1v_c) := 2, f'(x_1x_{p'}) := 4, f'(x_1u') := 5$, and $f'(x_{p'}v') := 5$. Then, extend f' to a (2,1)-total labeling of G according to Lemma 1.
 - 3.2. If $u' = v'$, then let $f'(x_1) := 0, f'(x_{p'}) := 2, f'(u') := 1, f'(x_1v_c) := 2, f'(x_1x_{p'}) := 4, f'(x_1u') := 3$, and $f'(x_{p'}u') := 5$.

□

4 Case $\Delta(G) = 4$

Let $G = (V, E)$ be an outerplane graph. The following structural properties of outerplane graphs are known.

Theorem 2. [14] *Every outerplane graph G with $\delta(G) = 2$ contains one of the following configurations:*

- (C1) two adjacent 2-vertices u_1 and u_2 .
- (C2) a 3-face $[u_1u_2u_3]$ with $d(u_1) = 2$ and $d(u_2) = 3$.
- (C3) two 3-faces $[u_1u_2u_3]$ and $[u_3u_4u_5]$ such that $d(u_3) = 4$ and $d(u_2) = d(u_4) = 2$.

In [4], Chen and Wang proved that if G does not contain (C3) and $\Delta(G) = 4$ holds, then $\lambda_2^T(G) \leq 6$. Their proof utilizes the property that if G contains (C1) or (C2), then a (2,1)-total labeling of G can be extended from a (2,1)-total labeling of some proper subgraph of G . Here, we investigate the case where G contains neither (C1) nor (C2), and derive a new structural property, i.e, if G contains neither (C1) nor (C2) and $\Delta(G) = 4$ holds, then G contains a new configuration as shown in Lemma 2 below, which includes (C3) as a subconfiguration.

Assume that $\Delta(G) = 4$. Define a *chain of 3-faces* as a sequence

$$C = \{[u_1u_2u_3], [u_3u_4u_5], \dots, [u_{2t-1}u_{2t}u_{2t+1}]\} \ (t \geq 2)$$

of 3-faces such that (u_{2i-1}, u_{2i}) and (u_{2i}, u_{2i+1}) are on the outer face of G and (u_{2i-1}, u_{2i+1}) is an inner edge for each $i = 1, 2, \dots, t$ (see Fig. 1). Notice that $d(u_{2i}) = 2$ holds for each $i = 1, 2, \dots, t$ and that $d(u_{2i+1}) = 4$ holds for




Fig. 1. A chain of 3-faces.

each $i = 1, 2, \dots, t - 1$ by $\Delta(G) = 4$; no vertex other than $u_{2i-1}, u_{2i}, u_{2i+2}$, and u_{2i+3} is adjacent to u_{2i+1} .

Lemma 2. *If G is an outerplane graph with $\Delta(G) = 4$ and $\delta(G) = 2$ which contains neither (C1) nor (C2), then G has a chain $\{[u_1u_2u_3], [u_3u_4u_5], \dots, [u_{2t-1}u_{2t}u_{2t+1}]\}$ of 3-faces such that (u_1, u_{2t+1}) is an inner edge.*

Proof. Let G be an outerplane graph with $\delta(G) = 2$ which contains neither (C1) nor (C2). Let G' be a 2-connected component of G which contains exactly one cut vertex v_c of G if G is not 2-connected, $G' = G$ otherwise. Notice that $|V(G')| \geq 3$ by $\delta(G) = 2$. Then G' has an inner edge since otherwise G contains (C1). There are the following two possible cases: (Case-I) every inner edge in G' belongs to some endface, (Case-II) otherwise. Assume that each vertex on the boundary of the outerface of G' is numbered $v'_1, v'_2, \dots, v'_t$ in clockwise order. Notice that each endface not containing v_c as a 2-vertex in G' is a 3-face since otherwise it would contain (C1).

(Case-I) Notice that there are at least two end faces in G' . Without loss of generality, let $[v'_1v'_2v'_3]$ be an endface of G' with $v_c \notin \{v'_2, v'_3\}$; $d_G(v'_2) = d_{G'}(v'_2) = 2$. Then an inner edge other than (v'_1, v'_3) is incident to v'_3 , since otherwise $d_G(v'_3) = 3$ holds and the 3-face $[v'_1v'_2v'_3]$ would satisfy the conditions of (C2). By the assumption of Case-I, the inner edge belongs to some endface, and hence we can see that another 3-face $[v'_3v'_4v'_5]$ exists; $\{[v'_1v'_2v'_3], [v'_3v'_4v'_5]\}$ is a chain. Moreover, observe that if $v'_1 \neq v_c$ also holds, then $[v'_{t-1}v'_tv'_1]$ is a 3-face and $\{[v'_{t-1}v'_tv'_1], [v'_1v'_2v'_3], [v'_3v'_4v'_5]\}$ is a chain. By repeating similar observations, it follows that there exists a chain $C = \{[v'_1v'_2v'_3], [v'_3v'_4v'_5], \dots, [v'_{t-1}v'_tv'_1]\}$. Then notice that $d_{G'}(v'_i) = 4$ (resp., 2) if i is odd (resp., even); by $\Delta(G) = 4$, if v_c exists, then v_c is some vertex v'_j with $d_{G'}(v'_j) = 2$. Hence, we can see that if v_c exists, then $G' - v_c$ contains a chain with an inner edge (v'_{j-1}, v'_{j+1}) , and that otherwise $C - \{[v'_{t-1}v'_tv'_1]\}$ is a chain with an inner edge (v'_1, v'_{t-1}) as required.

(Case-II) For an inner edge $(v'_i, v'_j) \in E$, let $G[v'_i, v'_j]$ denote the subgraph of G induced by $\{v'_i, v'_{i+1}, \dots, v'_{j-1}, v'_j\}$. Without loss of generality, let (v'_1, v'_i) be an inner edge in G' which does not belong to any endface such that $G[v'_2, v'_{i-1}]$ does not contain v_c . Moreover, we can choose such an edge (v'_1, v'_i) that any inner edge in $G[v'_1, v'_i]$ belongs to some endface of G . Note that if $G[v'_1, v'_i]$ contains an inner edge $e' = (v'_j, v'_k)$ ($1 \leq j < k \leq i$) not in any endface, then the number of inner edges not in any endface in $G[v'_j, v'_k]$ is less than that in $G[v'_1, v'_i]$ and this

observation indicates that this choice of (v'_1, v'_i) is possible. Then by applying the similar arguments in Case-I to endfaces in $G[v'_1, v'_i]$ (also in G), we can observe that there exists a chain $\{[v'_1 v'_2 v'_3], [v'_3 v'_4 v'_5], \dots, [v'_{i-2} v'_{i-1} v'_i]\}$, which is a required chain. $\square$

By utilizing this lemma, we show that $\lambda_2^T(G) \leq \Delta(G) + 2$ holds even if $\Delta(G) = 4$.

Theorem 3. *If $G = (V, E)$ is an outerplane graph with $\Delta(G) = 4$, then $\lambda_2^T(G) \leq 6$.*

Proof. We prove this by induction on $k = |V(G)| + |E(G)|$. The theorem clearly holds if $k \leq 9$. Consider the case of $k \geq 10$ and assume that for each $k' < k$, this theorem holds. We also assume that G is connected, since otherwise we can treat each component separately. Thus, $1 \leq \delta(G) \leq 2$.

In the case of $\delta(G) = 1$, similarly to the proof of Theorem 1, we can observe that G has a (2,1)-total labeling $f : V(G) \cup E(G) \rightarrow \mathcal{L}_6$.

Consider the case where $\delta(G) = 2$. By Theorem 2, G contains at least one of the configurations (C1)–(C3). In [4, Theorem 13], Chen and Wang showed that if G contains (C1) or (C2), then a (2,1)-total labeling of G can be obtained from a feasible labeling of some subgraph H of G . Here we only consider the case where G contains neither (C1) nor (C2). By Lemma 2, G contains a chain $C = \{[u_1 u_2 u_3], [u_3 u_4 u_5], \dots, [u_{2t-1} u_{2t} u_{2t+1}]\}$ of 3-faces such that (u_1, u_{2t+1}) is an inner edge. Let w_1 (resp., w_{2t+1}) denote the vertex not in C which is adjacent to u_1 (resp., u_{2t+1}) (note that by $\Delta(G) = 4$, no edge other than (u_1, w_1) and (u_{2t+1}, w_{2t+1}) connects C and the remaining part of G). By the induction hypothesis, $H := (G - \{u_2, u_3, \dots, u_{2t}\}) - \{(u_1, u_{2t+1})\}$ has a (2,1)-total labeling $f : V(H) \cup E(H) \rightarrow \mathcal{L}_6$. Then by $d_H(u_1) = d_H(u_{2t+1}) = 1$, we can assign any label in $\mathcal{L}_6 - \{f(w_1), f(u_1 w_1) - 1, f(u_1 w_1), f(u_1 w_1) + 1\}$ (resp., $\mathcal{L}_6 - \{f(w_{2t+1}), f(u_{2t+1} w_{2t+1}) - 1, f(u_{2t+1} w_{2t+1}), f(u_{2t+1} w_{2t+1}) + 1\}$) to u_1 (resp., u_{2t+1}) without violating the feasibility. Let $L(u_1; f)$ and $L(u_{2t+1}; f)$ denote the sets of such possible labels with respect to f for u_1 and u_{2t+1} , respectively; $L(u_1; f) = \mathcal{L}_6 - \{f(w_1), f(u_1 w_1) - 1, f(u_1 w_1), f(u_1 w_1) + 1\}$ and $L(u_{2t+1}; f) = \mathcal{L}_6 - \{f(w_{2t+1}), f(u_{2t+1} w_{2t+1}) - 1, f(u_{2t+1} w_{2t+1}), f(u_{2t+1} w_{2t+1}) + 1\}$. Notice that $|L(u_1; f)| \geq 3$ and $|L(u_{2t+1}; f)| \geq 3$.

Claim. $L(u_1; f) \times L(u_{2t+1}; f)$ contains one of $(0, 6), (0, 1), (1, 2), (2, 3), (3, 4), (4, 5), (5, 6), (6, 0), (6, 5), (5, 4), (4, 3), (3, 2), (2, 1)$, and $(1, 0)$.

Proof. Assume for contradiction that this claim does not hold. First consider the case of $\{0, 6\} \subseteq L(u_1; f)$. By assumption, $\{0, 6\} \cap L(u_{2t+1}; f) = \emptyset$. This indicates that (a1) $f(w_{2t+1}) = 0$ and $f(u_{2t+1} w_{2t+1}) \in \{5, 6\}$ or (a2) $f(w_{2t+1}) = 6$ and $f(u_{2t+1} w_{2t+1}) \in \{0, 1\}$. In the case of (a1), we have $1 \in L(u_{2t+1}; f)$ (i.e., $(0, 1) \in L(u_1; f) \times L(u_{2t+1}; f)$) and in the case of (a2), we have $5 \in L(u_{2t+1}; f)$ (i.e., $(6, 5) \in L(u_1; f) \times L(u_{2t+1}; f)$), a contradiction.

Next consider the case of $0 \in L(u_1; f)$ and $6 \notin L(u_1; f)$. By assumption, $\{1, 6\} \cap L(u_{2t+1}; f) = \emptyset$. This indicates that (b1) $f(w_{2t+1}) = 1$ and $f(u_{2t+1} w_{2t+1}) = 5$, (b2) $f(w_{2t+1}) = 1$ and $f(u_{2t+1} w_{2t+1}) = 6$, (b3) $f(w_{2t+1}) = 6$ and $f(u_{2t+1} w_{2t+1}) = 0$, (b4) $f(w_{2t+1}) = 6$ and $f(u_{2t+1} w_{2t+1}) = 1$, or (b5) $f(w_{2t+1}) = 6$ and $f(u_{2t+1} w_{2t+1}) = 2$. In the case of (b1) (resp., (b2), (b3), (b4), (b5)), we have $L(u_{2t+1}; f) = \{0, 2, 3\}$ (resp., $\{0, 2, 3, 4\}, \{2, 3, 4, 5\}, \{3, 4, 5\}, \{0, 4, 5\}$). By assumption that f does not satisfy this claim, it follows that in the case of (b1) (resp., (b2), (b3), (b4), (b5)), $L(u_1; f) \subseteq \{0, 5\}$ (resp., $\{0\}, \{0\}, \{0, 1\}, \{0, 2\}$). All of these cases contradict $|L(u_1; f)| \geq 3$.

By symmetry of labelings, the case of $6 \in L(u_1; f)$ and $0 \notin L(u_1; f)$ can be treated similarly to the previous case. Finally, consider the case of $\{0, 6\} \cap L(u_1; f) = \emptyset$. Assume that $\{0, 6\} \cap L(u_{2t+1}; f) = \emptyset$, since any other case can be treated similarly by exchanging parts of u_1 and u_{2t+1} . Then we can observe that $3 \in L(u_1; f)$, $\{2, 4\} \cap L(u_1; f) \neq \emptyset$, $3 \in L(u_{2t+1}; f)$, and $\{2, 4\} \cap L(u_{2t+1}; f) \neq \emptyset$. It follows that $(2, 3)$ or $(4, 3)$ are contained in $L(u_1; f) \times L(u_{2t+1}; f)$, a contradiction. $\square$

By symmetry, it suffices to consider the following four cases: (Case-1) $0 \in L(u_1; f)$ and $6 \in L(u_{2t+1}; f)$, (Case-2) $0 \in L(u_1; f)$ and $1 \in L(u_{2t+1}; f)$, (Case-3) $1 \in L(u_1; f)$ and $2 \in L(u_{2t+1}; f)$, and (Case-4) $2 \in L(u_1; f)$ and $3 \in L(u_{2t+1}; f)$. In each case, we will extend f to a (2,1)-total labeling $f' : V \cup E \rightarrow \mathcal{L}_6$ of G . Let $f'(x) := f(x)$ for every $x \in V(H) \cup E(H)$.

(Case-1) Let $f'(u_1) := 0$ and $f'(u_{2t+1}) := 6$.

(1.1) Assume that t is even; $t = 2k$.

(1.1.1) Assume that $f'(u_1 w_1) \neq 6$ and $f'(u_{2t+1} w_{2t+1}) \neq 0$.

Let $f'(u_2) := 6, f'(u_3) := 3, f'(u_4) := 0, f'(u_5) := 6, f'(u_2 u_3) := 1, f'(u_1 u_3) := 6, f'(u_3 u_4) := 5, f'(u_4 u_5) := 4$, and $f'(u_3 u_5) := 0$. For $i = 2, 3, \dots, k$, let $f'(u_{4i-2}) := 1, f'(u_{4i-1}) := 3, f'(u_{4i}) := 0, f'(u_{4i+1}) := 6, f'(u_{4i-3} u_{4i-2}) := 3, f'(u_{4i-2} u_{4i-1}) := 6, f'(u_{4i-3} u_{4i-1}) := 1, f'(u_{4i-1} u_{4i}) := 5, f'(u_{4i} u_{4i+1}) := 4$, and $f'(u_{4i-1} u_{4i+1}) := 0$. Denote this labeling f' by f_1 (see Fig. 2). We next assign a label $\{2, 3, 4\} - \{f'(u_1 w_1), f'(u_{2t+1} w_{2t+1})\}$ to (u_1, u_{2t+1}) , and a label in $\{2, 3, 4\} - \{f'(u_1 w_1), f'(u_{2t+1} w_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{2, 3, 4\} - \{f'(u_{2t+1} w_{2t+1}), f'(u_1 u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then the resulting labeling f' is a (2,1)-total labeling of G .




Fig. 2. A labeling f_1 on C .

(1.1.2) Assume that $f'(u_1w_1) = 6$ and $f'(u_{2t+1}w_{2t+1}) \neq 0$.

Let $f' := f_1$. We reassign labels for some vertices and edges as follows. Let $f'(u_2) := 1$, $f'(u_3) := 2$, $f'(u_1u_2) := 5$, $f'(u_2u_3) := 6$ and $f'(u_1u_3) := 4$. Next we assign a label in $\{2, 3\} - \{f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) and reassign a label in $\{2, 3, 4\} - \{f'(u_1u_{2t+1}), f'(u_{2t+1}w_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then the resulting labeling f' is a $(2,1)$ -total labeling of G .

(1.1.3) Assume that $f'(u_1w_1) \neq 6$ and $f'(u_{2t+1}w_{2t+1}) = 0$.

Let $f' := f_1$. We reassign labels for some vertices and edges as follows. Let $f'(u_{2t-1}) := 4$, $f'(u_{2t}) := 5$, $f'(u_{2t-1}u_{2t}) := 0$, $f'(u_{2t}u_{2t+1}) := 1$, and $f'(u_{2t-1}u_{2t+1}) := 2$. Next we assign a label in $\{3, 4\} - \{f'(u_1w_1)\}$ to (u_1, u_{2t+1}) and a label in $\{2, 3, 4\} - \{f'(u_1u_{2t+1}), f'(u_1w_1)\}$ to (u_1, u_2) . Then the resulting labeling f' is a $(2,1)$ -total labeling of G .

(1.1.4) Assume that $f'(u_1w_1) = 6$ and $f'(u_{2t+1}w_{2t+1}) = 0$.

We can obtain a $(2,1)$ -total labeling f' of G as follows. If $t = 2$, then let $f'(u_2) := 3$, $f'(u_3) := 1$, $f'(u_4) := 3$, $f'(u_1u_2) := 5$, $f'(u_2u_3) := 6$, $f'(u_1u_3) := 4$, $f'(u_3u_4) := 5$, $f'(u_4u_5) := 1$, $f'(u_3u_5) := 3$, and $f'(u_1u_5) := 2$.

Assume that $t \geq 4$. Let $f' := f_1$. We reassign labels for some vertices and edges similarly to (1.1.2) and (1.1.3). Namely, let $f'(u_2) := 1$, $f'(u_3) := 2$, $f'(u_1u_2) := 5$, $f'(u_2u_3) := 6$, $f'(u_1u_3) := 4$, $f'(u_{2t-1}) := 4$, $f'(u_{2t}) := 5$, $f'(u_{2t-1}u_{2t}) := 0$, $f'(u_{2t}u_{2t+1}) := 1$, $f'(u_{2t-1}u_{2t+1}) := 2$, and $f'(u_1u_{2t+1}) := 3$.

(1.2) Assume that t is odd; $t = 2k + 1$ ($k \geq 1$).

(1.2.1) Assume that $f'(u_1w_1) \neq 6$ and $f'(u_{2t+1}w_{2t+1}) \neq 0$.

Let $f'(u_2) := 6$, $f'(u_3) := 3$, $f'(u_4) := 0$, $f'(u_5) := 4$, $f'(u_6) := 0$, $f'(u_7) := 6$, $f'(u_2u_3) := 0$, $f'(u_1u_3) := 6$, $f'(u_3u_4) := 5$, $f'(u_4u_5) := 2$, $f'(u_3u_5) := 1$, $f'(u_5u_6) := 6$, $f'(u_6u_7) := 4$, and $f'(u_5u_7) := 0$. For $i = 2, 3, \dots, k$, let $f'(u_{4i}) := 0$, $f'(u_{4i+1}) := 4$, $f'(u_{4i+2}) := 0$, $f'(u_{4i+3}) := 6$, $f'(u_{4i-1}u_{4i}) := 3$, $f'(u_{4i}u_{4i+1}) := 2$, $f'(u_{4i-1}u_{4i+1}) := 1$, $f'(u_{4i+1}u_{4i+2}) := 6$, $f'(u_{4i+2}u_{4i+3}) := 4$, and $f'(u_{4i+1}u_{4i+3}) := 0$. Denote this labeling f' by f_2 (see Fig. 3). We next assign a label $\{2, 3, 4\} - \{f'(u_1w_1), f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) and assign a label in $\{2, 3, 4\} - \{f'(u_1w_1), f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{2, 3, 4\} - \{f'(u_{2t+1}w_{2t+1}), f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) .

(1.2.2) Assume that $f'(u_1w_1) = 6$ and $f'(u_{2t+1}w_{2t+1}) \neq 0$.

Let $f' := f_2$. We reassign labels for some vertices and edges as follows. Let $f'(u_2) := 2$, $f'(u_3) := 6$, $f'(u_1u_2) := 5$, $f'(u_1u_3) := 4$, and $f'(u_3u_4) := 3$. Next we assign a label in $\{2, 3\} - \{f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) . Reassign a label in $\{2, 3, 4\} - \{f'(u_1u_{2t+1}), f'(u_{2t+1}w_{2t+1})\}$ to (u_{2t}, u_{2t+1}) .

(1.2.3) Assume that $f'(u_1w_1) \neq 6$ and $f'(u_{2t+1}w_{2t+1}) = 0$.

Let $f' := f_2$. We reassign labels for some vertices and edges as follows. Let $f'(u_{2t-1}) := 5$, $f'(u_{2t}) := 3$, $f'(u_{2t-1}u_{2t}) := 0$, $f'(u_{2t}u_{2t+1}) := 1$, and $f'(u_{2t-1}u_{2t+1}) := 3$. Next we assign a label in $\{2, 4\} - \{f'(u_1w_1)\}$ to (u_1, u_{2t+1}) and a label in $\{2, 3, 4\} - \{f'(u_1w_1), f'(u_1u_{2t+1})\}$ to (u_1, u_2) .

(1.2.4) Assume that $f'(u_1w_1) = 6$ and $f'(u_{2t+1}w_{2t+1}) = 0$.




Fig. 3. A labeling f_2 on C .

Let $f' := f_2$. We reassign labels for some vertices and edges similarly to (1.2.2) and (1.2.3). Namely, let $f'(u_2) := 2$, $f'(u_3) := 6$, $f'(u_1u_2) := 5$, $f'(u_1u_3) := 4$, $f'(u_3u_4) := 3$, $f'(u_{2t-1}) := 5$, $f'(u_{2t}) := 3$, $f'(u_{2t-1}u_{2t}) := 0$, $f'(u_{2t}u_{2t+1}) := 1$, $f'(u_{2t-1}u_{2t+1}) := 3$, and $f'(u_1u_{2t+1}) := 2$.

(Case-2) Let $f'(u_1) := 0$ and $f'(u_{2t+1}) := 1$.

(2.1) Assume that t is even; $t = 2k$.

(2.1.1) Assume that $f'(u_1w_1) \neq 2$ and $f'(u_{2t+1}w_{2t+1}) \notin \{3, 4\}$.

Let $f'(u_2) := 2$, $f'(u_3) := 6$, $f'(u_4) := 5$, $f'(u_5) := 1$, $f'(u_2u_3) := 0$, $f'(u_1u_3) := 2$, $f'(u_3u_4) := 1$, $f'(u_4u_5) := 3$, and $f'(u_3u_5) := 4$. For $i = 2, 3, \dots, k$, let $f'(u_{4i-2}) := 4$, $f'(u_{4i-1}) := 0$, $f'(u_{4i}) := 2$, $f'(u_{4i+1}) := 1$, $f'(u_{4i-3}u_{4i-2}) := 6$, $f'(u_{4i-2}u_{4i-1}) := 2$, $f'(u_{4i-3}u_{4i-1}) := 5$, $f'(u_{4i-1}u_{4i}) := 6$, $f'(u_{4i}u_{4i+1}) := 4$, and $f'(u_{4i-1}u_{4i+1}) := 3$. Denote this labeling f' by f_3 (see Fig. 4).




Fig. 4. A labeling f_3 on C .

Consider the case of $t = 2$. We assign a label $\{3, 5, 6\} - \{f'(u_1w_1), f'(u_5w_5)\}$ to (u_1, u_5) and a label in $\{4, 5, 6\} - \{f'(u_1w_1), f'(u_1u_5)\}$ to (u_1, u_2) . Reassign a label in $\{3, 5, 6\} - \{f'(u_5w_5), f'(u_1u_5)\}$ to (u_4, u_5) . Then if $f'(u_4u_5) \in \{5, 6\}$, then reassign a label for u_4 as $f'(u_4) := 3$.

Consider the case of $t \geq 4$. Assign a label $\{4, 5, 6\} - \{f'(u_1w_1), f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) and a label in $\{4, 5, 6\} - \{f'(u_1w_1), f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{4, 5, 6\} - \{f'(u_{2t+1}w_{2t+1}), f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_{2t}u_{2t+1}) = 6$, then reassign a label for (u_{2t-1}, u_{2t}) as $f'(u_{2t-1}u_{2t}) := 4$.

(2.1.2) Assume that $f'(u_1w_1) = 2$ and $f'(u_{2t+1}w_{2t+1}) \notin \{3, 4\}$.

Let $f' := f_3$. We reassign a label for (u_1, u_3) to 3.

Consider the case of $t = 2$. Assign a label in $\{5, 6\} - \{f'(u_5w_5)\}$ to (u_1, u_5) and a label in $\{4, 5, 6\} - \{f'(u_1u_5)\}$ to (u_1, u_2) . Reassign a label in $\{3, 5, 6\} - \{f'(u_5w_5), f'(u_1u_5)\}$ to (u_4, u_5) . Then if $f'(u_4u_5) \in \{5, 6\}$, then let $f'(u_4) := 3$.

Consider the case of $t \geq 4$. Assign a label in $\{4, 5, 6\} - \{f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) and a label in $\{4, 5, 6\} - \{f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{4, 5, 6\} - \{f'(u_{2t+1}w_{2t+1}), f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then, if $f'(u_{2t}u_{2t+1}) = 6$, then let $f'(u_{2t-1}u_{2t}) := 4$.

(2.1.3) Assume that $f'(u_1w_1) \neq 2$ and $f'(u_{2t+1}w_{2t+1}) = 3$.

If $t = 2$, then we can obtain a feasible labeling similarly to (2.1.1). Consider the case of $t \geq 4$. Let $f' := f_3$. Reassign a label for (u_{2t-1}, u_{2t+1}) as $f'(u_{2t-1}u_{2t+1}) := 6$. Next we assign a label in $\{4, 5\} - \{f'(u_1w_1)\}$ to (u_1, u_{2t+1}) and a label in $\{4, 5, 6\} - \{f'(u_1u_{2t+1}), f'(u_1w_1)\}$ to (u_1, u_2) . Reassign a label in $\{4, 5\} - \{f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_{2t}u_{2t+1}) = 4$, then let $f'(u_{2t}) := 6$ and $f'(u_{2t-1}u_{2t}) := 3$, and if $f'(u_{2t}u_{2t+1}) = 5$, then let $f'(u_{2t-1}u_{2t}) := 4$.

(2.1.4) Assume that $f'(u_1w_1) \neq 2$ and $f'(u_{2t+1}w_{2t+1}) = 4$.

If $t \geq 4$, then we can obtain a feasible labeling similarly to (2.1.1). Consider the case of $t = 2$. Let $f' := f_3$. We reassign labels for (u_3, u_5) and u_4 as $f'(u_3u_5) := 3$ and $f'(u_4) := 3$. Next we assign a label in $\{5, 6\} - \{f'(u_1w_1)\}$ to (u_1, u_5) and a label in $\{4, 5, 6\} - \{f'(u_1u_5), f'(u_1w_1)\}$ to (u_1, u_2) . Reassign a label in $\{5, 6\} - \{f'(u_1u_5)\}$ to (u_4, u_5) .

(2.1.5) Assume that $f'(u_1w_1) = 2$ and $f'(u_{2t+1}w_{2t+1}) = 3$.

If $t = 2$, then we can obtain a feasible labeling similarly to (2.1.2). Consider the case of $t \geq 4$. Let $f' := f_3$. We reassign labels for some vertices and edges similarly to (2.1.2) and (2.1.3). Namely, let $f'(u_1u_2) := 4$, $f'(u_1u_3) := 3$, $f'(u_{2t}) := 6$, $f'(u_{2t-1}u_{2t}) := 3$, $f'(u_{2t}u_{2t+1}) := 4$, $f'(u_{2t-1}u_{2t+1}) := 6$, and $f'(u_1u_{2t+1}) := 5$.

(2.1.6) Assume that $f'(u_1w_1) = 2$ and $f'(u_{2t+1}w_{2t+1}) = 4$.

If $t \geq 4$, then we can obtain a feasible labeling similarly to (2.1.2). If $t = 2$, let $f'(u_2) := 2$, $f'(u_3) := 6$, $f'(u_4) := 3$, $f'(u_1u_2) := 5$, $f'(u_2u_3) := 0$, $f'(u_1u_3) := 4$, $f'(u_3u_4) := 1$, $f'(u_4u_5) := 5$, $f'(u_3u_5) = 3$, and $f'(u_1u_5) := 6$.

(2.2) Assume that t is odd; $t = 2k + 1$ ($k \geq 1$).

(2.2.1) Assume that $f'(u_1w_1) \neq 2$ and $f'(u_{2t+1}w_{2t+1}) \neq 5$.

Let $f'(u_2) := 2$, $f'(u_3) := 6$, $f'(u_4) := 4$, $f'(u_5) := 0$, $f'(u_6) := 6$, $f'(u_7) := 1$, $f'(u_2u_3) := 0$, $f'(u_1u_3) := 2$, $f'(u_3u_4) := 1$, $f'(u_4u_5) := 6$, $f'(u_3u_5) := 4$, $f'(u_5u_6) := 2$, $f'(u_6u_7) := 3$, and $f'(u_5u_7) := 5$. For $i = 2, 3, \dots, k$, let $f'(u_{4i}) := 6$, $f'(u_{4i+1}) := 0$, $f'(u_{4i+2}) := 6$, $f'(u_{4i+3}) := 1$, $f'(u_{4i-1}u_{4i}) := 4$, $f'(u_{4i}u_{4i+1}) := 3$, $f'(u_{4i-1}u_{4i+1}) := 6$, $f'(u_{4i+1}u_{4i+2}) := 2$, $f'(u_{4i+2}u_{4i+3}) := 3$, and $f'(u_{4i+1}u_{4i+3}) := 5$. Denote this labeling f' by f_4 (see Fig. 5). We next assign a label $\{3, 4, 6\} - \{f'(u_1w_1), f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) . and a label in $\{4, 5, 6\} - \{f'(u_1w_1), f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{3, 4, 6\} - \{f'(u_{2t+1}w_{2t+1}), f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_{2t}u_{2t+1}) = 6$, then let $f'(u_{2t}) := 4$.

(2.2.2) Assume that $f'(u_1w_1) = 2$ and $f'(u_{2t+1}w_{2t+1}) \neq 5$.

Let $f' := f_4$. We reassign labels for some vertices and edges as follows. Let $f'(u_1u_2) := 5$ and $f'(u_1u_3) := 3$. Next we assign a label in $\{4, 6\} - \{f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) . Reassign a label in $\{3, 4, 6\} - \{f'(u_1u_{2t+1}), f'(u_{2t+1}w_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_{2t}u_{2t+1}) = 6$, then let $f'(u_{2t}) := 4$.

(2.2.3) Assume that $f'(u_1w_1) \neq 2$ and $f'(u_{2t+1}w_{2t+1}) = 5$.

Let $f' := f_4$.

First consider the case of $t = 3$. Let $f'(u_6) := 2$, $f'(u_5u_6) := 5$, and $f'(u_5u_7) := 3$. Next we assign a label in $\{4, 6\} - \{f'(u_1w_1)\}$ to (u_1, u_7) and a label in $\{4, 5, 6\} - \{f'(u_1w_1), f'(u_1u_7)\}$ to (u_1, u_2) . Reassign a label in $\{4, 6\} - \{f'(u_1u_7)\}$ to (u_6, u_7) .

Next consider the case of $t \geq 5$. Let $f'(u_{2t-1}u_{2t+1}) := 4$. Next we assign a label in $\{3, 6\} - \{f'(u_1w_1)\}$ to (u_1, u_{2t+1}) and a label in $\{4, 5, 6\} - \{f'(u_1w_1), f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{3, 6\} - \{f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_{2t}u_{2t+1}) = 6$, then $f'(u_{2t}) := 4$.

(2.2.4) Assume that $f'(u_1w_1) = 2$ and $f'(u_{2t+1}w_{2t+1}) = 5$.

Let $f' := f_4$. We reassign labels for some vertices and edges similarly to (2.2.2) and (2.2.3). Namely, if $t = 3$, let $f'(u_1u_2) := 5$, $f'(u_1u_3) := 3$, $f'(u_6) := 2$, $f'(u_5u_6) := 5$, $f'(u_6u_7) := 6$, $f'(u_5u_7) := 3$, and $f'(u_1u_7) := 4$. If $t \geq 5$, then let $f'(u_1u_2) := 5$, $f'(u_1u_3) := 3$, $f'(u_{2t-1}u_{2t+1}) := 4$, and $f'(u_1u_{2t+1}) := 6$.

(Case-3) Let $f'(u_1) := 1$ and $f'(u_{2t+1}) := 2$.




Fig. 5. A labeling f_4 on C .

(3.1) Assume that t is even; $t = 2k$.

(3.1.1) Assume that $f'(u_1w_1) \neq 3$ and $f'(u_{2t+1}w_{2t+1}) \neq 0$.

Let $f'(u_2) := 0$, $f'(u_3) := 5$, $f'(u_4) := 3$, $f'(u_5) := 2$, $f'(u_2u_3) := 2$, $f'(u_1u_3) := 3$, $f'(u_3u_4) := 1$, $f'(u_4u_5) := 6$, and $f'(u_3u_5) := 0$. For $i = 2, 3, \dots, k$, let $f'(u_{4i-2}) := 0$, $f'(u_{4i-1}) := 6$, $f'(u_{4i}) := 0$, $f'(u_{4i+1}) := 2$, $f'(u_{4i-3}u_{4i-2}) := 5$, $f'(u_{4i-2}u_{4i-1}) := 3$, $f'(u_{4i-3}u_{4i-1}) := 4$, $f'(u_{4i-1}u_{4i}) := 2$, $f'(u_{4i}u_{4i+1}) := 6$, and $f'(u_{4i-1}u_{4i+1}) := 0$. Denote this labeling f' by f_5 (see Fig. 6). Assign a label $\{4, 5, 6\} - \{f'(u_1w_1), f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) and




Fig. 6. A labeling f_5 on C .

a label in $\{4, 5, 6\} - \{f'(u_1w_1), f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{4, 5, 6\} - \{f'(u_{2t+1}w_{2t+1}), f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_4u_5) = 4$, then let $f'(u_4) := 6$.

(3.1.2) Assume that $f'(u_1w_1) = 3$ and $f'(u_{2t+1}w_{2t+1}) \neq 0$.

Let $f' := f_5$. We reassign labels for some vertices and edges as follows. Let $f'(u_3) := 6$, $f'(u_2u_3) := 3$, and $f'(u_1u_3) := 4$. Assign a label in $\{5, 6\} - \{f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) and a label in $\{5, 6\} - \{f'(u_1u_{2t+1})\}$ to (u_1, u_2) .

Reassign a label in $\{4, 5, 6\} - \{f'(u_{2t+1}w_{2t+1}), f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_4u_5) = 4$, then let $f'(u_4) := 0$ and $f'(u_3u_4) := 2$.

(3.1.3) Assume that $f'(u_1w_1) \neq 3$ and $f'(u_{2t+1}w_{2t+1}) = 0$.

Let $f' := f_5$.

Consider the case of $t = 2$. Let $f'(u_3u_5) := 4$ and $f'(u_3) := 6$. Next we assign a label in $\{5, 6\} - \{f'(u_1w_1)\}$ to (u_1, u_5) and a label in $\{4, 5, 6\} - \{f'(u_1u_5), f'(u_1w_1)\}$ to (u_1, u_2) . Reassign a label in $\{5, 6\} - \{f'(u_1u_5)\}$ to (u_4, u_5) .

Consider the case of $t \geq 4$. Let $f'(u_{2t-1}u_{2t+1}) := 5$, $f'(u_{2t-2}) := 1$, and $f'(u_{2t-1}) := 0$. Next we assign a label in $\{4, 6\} - \{f'(u_1w_1)\}$ to (u_1, u_{2t+1}) and a label in $\{4, 5, 6\} - \{f'(u_1u_{2t+1}), f'(u_1w_1)\}$ to (u_1, u_2) . Reassign a label in $\{4, 6\} - \{f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_{2t}u_{2t+1}) = 4$, then let $f'(u_{2t}) := 6$, and if $f'(u_{2t}u_{2t+1}) = 6$, then let $f'(u_{2t}) := 4$.

(3.1.4) Assume that $f'(u_1w_1) = 3$ and $f'(u_{2t+1}w_{2t+1}) = 0$.

If $t = 2$, let $f'(u_2) := 2$, $f'(u_3) := 0$, $f'(u_4) := 6$, $f'(u_1u_2) := 6$, $f'(u_2u_3) := 5$, $f'(u_1u_3) := 4$, $f'(u_3u_4) := 2$, $f'(u_4u_5) := 4$, $f'(u_3u_5) = 6$, and $f'(u_1u_5) := 5$.

Consider the case of $t \geq 4$. Let $f' := f_5$. We reassign labels for some vertices and edges similarly to (3.1.2) and (3.1.3). Namely, let $f'(u_1u_2) := 5$, $f'(u_3) := 6$, $f'(u_1u_3) := 4$, $f'(u_{2t-2}) := 1$, $f'(u_{2t-1}) := 0$, $f'(u_{2t}) := 6$, $f'(u_{2t}u_{2t+1}) := 4$, $f'(u_{2t-1}u_{2t+1}) := 5$, and $f'(u_1u_{2t+1}) := 6$.

(3.2) Assume that t is odd; $t = 2k + 1$ ($k \geq 1$).

(3.2.1) Assume that $f'(u_1w_1) \neq 3$ and $f'(u_{2t+1}w_{2t+1}) \neq 0$.

Let $f'(u_2) := 0$, $f'(u_3) := 5$, $f'(u_4) := 2$, $f'(u_5) := 6$, $f'(u_6) := 0$, $f'(u_7) := 2$, $f'(u_2u_3) := 2$, $f'(u_1u_3) := 3$, $f'(u_3u_4) := 0$, $f'(u_4u_5) := 4$, $f'(u_3u_5) := 1$, $f'(u_5u_6) := 2$, $f'(u_6u_7) := 6$, and $f'(u_5u_7) := 0$. For $i = 2, 3, \dots, k$, let $f'(u_{4i}) := 0$, $f'(u_{4i+1}) := 6$, $f'(u_{4i+2}) := 0$, $f'(u_{4i+3}) := 2$, $f'(u_{4i-1}u_{4i}) := 5$, $f'(u_{4i}u_{4i+1}) := 3$, $f'(u_{4i-1}u_{4i+1}) := 4$, $f'(u_{4i+1}u_{4i+2}) := 2$, $f'(u_{4i+2}u_{4i+3}) := 6$, and $f'(u_{4i+1}u_{4i+3}) := 0$. Denote this labeling f' by f_6 (see Fig. 7). We next




Fig. 7. A labeling f_6 on C .

assign a label $\{4, 5, 6\} - \{f'(u_1w_1), f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) and a label in $\{4, 5, 6\} - \{f'(u_1w_1), f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{4, 5, 6\} - \{f'(u_{2t+1}w_{2t+1}), f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) .

(3.2.2) Assume that $f'(u_1w_1) = 3$ and $f'(u_{2t+1}w_{2t+1}) \neq 0$.

Let $f' := f_6$. We reassign labels for some vertices and edges as follows. Let $f'(u_1u_3) := 6$ and $f'(u_3) := 4$. Next we assign a label in $\{4, 5\} - \{f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) and a label in $\{4, 5\} - \{f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{4, 5, 6\} - \{f'(u_1u_{2t+1}), f'(u_{2t+1}w_{2t+1})\}$ to (u_{2t}, u_{2t+1}) .

(3.2.3) Assume that $f'(u_1w_1) \neq 3$ and $f'(u_{2t+1}w_{2t+1}) = 0$.

Let $f' := f_6$.

First consider the case of $t = 3$. Let $f'(u_4) := 4$, $f'(u_4u_5) := 2$, $f'(u_5u_6) := 3$, and $f'(u_5u_7) := 4$. Next we assign a label in $\{5, 6\} - \{f'(u_1w_1)\}$ to (u_1, u_7) and a label in $\{4, 5, 6\} - \{f'(u_1w_1), f'(u_1u_7)\}$ to (u_1, u_2) . Reassign a label in $\{5, 6\} - \{f'(u_1u_7)\}$ to (u_6, u_7) .

Next consider the case of $t \geq 5$. Let $f'(u_{2t-2}) := 6$, $f'(u_{2t-1}) := 0$, $f'(u_{2t-3}u_{2t-2}) := 4$, $f'(u_{2t-3}u_{2t-1}) := 5$, and $f'(u_{2t-1}u_{2t+1}) := 6$. Next we assign a label in $\{4, 5\} - \{f'(u_1w_1)\}$ to (u_1, u_{2t+1}) and a label in $\{4, 5, 6\} - \{f'(u_1w_1), f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{4, 5\} - \{f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_{2t}u_{2t+1}) = 4$ (resp., 5), then let $f'(u_{2t}) := 6$ (resp., 1) and $f'(u_{2t-1}u_{2t}) := 2$ (resp., 4).

(3.2.4) Assume that $f'(u_1w_1) = 3$ and $f'(u_{2t+1}w_{2t+1}) = 0$.

Let $f' := f_6$. We reassign labels for some vertices and edges similarly to (3.2.2) and (3.2.3). Namely, if $t = 3$, let $f'(u_1u_2) := 4$, $f'(u_1u_3) := 6$, $f'(u_3) := 4$, $f'(u_4) := 5$, $f'(u_3u_4) := 1$, $f'(u_4u_5) := 2$, $f'(u_3u_5) := 0$, $f'(u_5u_6) := 3$, $f'(u_6u_7) := 6$, $f'(u_5u_7) := 4$, and $f'(u_1u_7) := 5$. If $t \geq 5$, then let $f'(u_1u_2) := 5$, $f'(u_1u_3) := 6$, $f'(u_3) := 4$, $f'(u_{2t-2}) := 6$, $f'(u_{2t-1}) := 0$, $f'(u_{2t}) := 1$, $f'(u_{2t-3}u_{2t-2}) := 4$, $f'(u_{2t-3}u_{2t-1}) := 5$, $f'(u_{2t-1}u_{2t}) := 4$, $f'(u_{2t}u_{2t+1}) := 5$, $f'(u_{2t-1}u_{2t+1}) := 6$, and $f'(u_1u_{2t+1}) := 4$.

(Case-4) Let $f'(u_1) := 2$ and $f'(u_{2t+1}) := 3$.

(4.1) Assume that t is even; $t = 2k$.

(4.1.1) Assume that $f'(u_1w_1) \neq 4$ and $f'(u_{2t+1}w_{2t+1}) \neq 1$.

Let $f'(u_2) := 1$, $f'(u_3) := 6$, $f'(u_4) := 4$, $f'(u_5) := 3$, $f'(u_2u_3) := 3$, $f'(u_1u_3) := 4$, $f'(u_3u_4) := 2$, $f'(u_4u_5) := 0$, and $f'(u_3u_5) := 1$. For $i = 2, 3, \dots, k$, let $f'(u_{4i-2}) := 2$, $f'(u_{4i-1}) := 4$, $f'(u_{4i}) := 5$, $f'(u_{4i+1}) := 3$, $f'(u_{4i-3}u_{4i-2}) := 5$, $f'(u_{4i-2}u_{4i-1}) := 0$, $f'(u_{4i-3}u_{4i-1}) := 6$, $f'(u_{4i-1}u_{4i}) := 2$, $f'(u_{4i}u_{4i+1}) := 0$, and $f'(u_{4i-1}u_{4i+1}) := 1$. Denote this labeling f' by f_7 (see Fig. 8). Assign a label $\{0, 5, 6\} - \{f'(u_1w_1), f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) and




Fig. 8. A labeling f_7 on C .

a label in $\{0, 5, 6\} - \{f'(u_1w_1), f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{0, 5, 6\} - \{f'(u_{2t+1}w_{2t+1}), f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_1u_2) = 0$, then let $f'(u_2) := 5$, and if $f'(u_{2t}u_{2t+1}) \in \{5, 6\}$, then let $f'(u_{2t}) := 0$.

(4.1.2) Assume that $f'(u_1w_1) = 4$ and $f'(u_{2t+1}w_{2t+1}) \neq 1$.

Let $f' := f_7$. We reassign a label for (u_1, u_3) as $f'(u_1u_3) := 0$. Assign a label in $\{5, 6\} - \{f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) and a label in $\{5, 6\} - \{f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{0, 5, 6\} - \{f'(u_{2t+1}w_{2t+1}), f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_{2t}u_{2t+1}) \in \{5, 6\}$, then let $f'(u_{2t}) := 0$.

(4.1.3) Assume that $f'(u_1w_1) \neq 4$ and $f'(u_{2t+1}w_{2t+1}) = 1$.

Let $f' := f_7$.

Consider the case of $t = 2$. Let $f'(u_3u_5) := 0$ and $f'(u_4) := 0$. Next we assign a label in $\{5, 6\} - \{f'(u_1w_1)\}$ to (u_1, u_5) and a label in $\{0, 5, 6\} - \{f'(u_1u_5), f'(u_1w_1)\}$ to (u_1, u_2) . Reassign a label in $\{5, 6\} - \{f'(u_1u_5)\}$ to (u_4, u_5) . Then if $f'(u_1u_2) = 0$, then let $f'(u_2) := 5$.

Consider the case of $t \geq 4$. Let $f'(u_{2t-1}u_{2t+1}) := 5$, $f'(u_{2t-2}) := 1$, $f'(u_{2t-1}) := 0$, $f'(u_{2t-2}u_{2t-1}) := 3$, and $f'(u_{2t}) := 4$. Next we assign a label in $\{0, 6\} - \{f'(u_1w_1)\}$ to (u_1, u_{2t+1}) and a label in $\{0, 5, 6\} - \{f'(u_1u_{2t+1}), f'(u_1w_1)\}$ to (u_1, u_2) . Reassign a label in $\{0, 6\} - \{f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_1u_2) = 0$, then let $f'(u_2) := 5$.

(4.1.4) Assume that $f'(u_1w_1) = 4$ and $f'(u_{2t+1}w_{2t+1}) = 1$.

If $t = 2$, let $f'(u_2) := 4$, $f'(u_3) := 0$, $f'(u_4) := 1$, $f'(u_1u_2) := 6$, $f'(u_2u_3) := 2$, $f'(u_1u_3) := 5$, $f'(u_3u_4) := 3$, $f'(u_4u_5) := 5$, $f'(u_3u_5) := 6$, and $f'(u_1u_5) := 0$.

Consider the case of $t \geq 4$. Let $f' := f_7$. We reassign labels for some vertices and edges similarly to (4.1.2) and (4.1.3). Namely, let $f'(u_1u_2) := 5$, $f'(u_1u_3) := 0$, $f'(u_{2t-2}) := 1$, $f'(u_{2t-1}) := 0$, $f'(u_{2t}) := 4$, $f'(u_{2t-2}u_{2t-1}) := 3$, $f'(u_{2t-1}u_{2t+1}) := 5$, and $f'(u_1u_{2t+1}) := 6$.




Fig. 9. A labeling f_8 on C .

(4.2) Assume that t is odd; $t = 2k + 1$ ($k \geq 1$).

(4.2.1) Assume that $f'(u_1w_1) \neq 4$ and $f'(u_{2t+1}w_{2t+1}) \neq 1$.

Let $f'(u_2) := 1$, $f'(u_3) := 0$, $f'(u_4) := 1$, $f'(u_5) := 6$, $f'(u_6) := 2$, $f'(u_7) := 3$, $f'(u_2u_3) := 3$, $f'(u_1u_3) := 4$, $f'(u_3u_4) := 5$, $f'(u_4u_5) := 3$, $f'(u_3u_5) := 2$, $f'(u_5u_6) := 4$, $f'(u_6u_7) := 0$, and $f'(u_5u_7) := 1$. For $i = 2, 3, \dots, k$, let $f'(u_{4i}) := 2$, $f'(u_{4i+1}) := 4$, $f'(u_{4i+2}) := 5$, $f'(u_{4i+3}) := 3$, $f'(u_{4i-1}u_{4i}) := 5$, $f'(u_{4i}u_{4i+1}) := 0$, $f'(u_{4i-1}u_{4i+1}) := 6$, $f'(u_{4i+1}u_{4i+2}) := 2$, $f'(u_{4i+2}u_{4i+3}) := 0$, and $f'(u_{4i+1}u_{4i+3}) := 1$. Denote this labeling f' by f_8 (see Fig. 9). We next assign a label $\{0, 5, 6\} - \{f'(u_1w_1), f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) and a label in $\{0, 5, 6\} - \{f'(u_1w_1), f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{0, 5, 6\} - \{f'(u_{2t+1}w_{2t+1}), f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_1u_2) = 0$, then let $f'(u_2) := 5$, and if $f'(u_{2t}u_{2t+1}) \in \{5, 6\}$, then let $f'(u_{2t}) := 0$.

(4.2.2) Assume that $f'(u_1w_1) = 4$ and $f'(u_{2t+1}w_{2t+1}) \neq 1$.

Let $f' := f_8$. We reassign a label for (u_1, u_3) as $f'(u_1u_3) := 6$. Next we assign a label in $\{0, 5\} - \{f'(u_{2t+1}w_{2t+1})\}$ to (u_1, u_{2t+1}) and a label in $\{0, 5\} - \{f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{0, 5, 6\} - \{f'(u_1u_{2t+1}), f'(u_{2t+1}w_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_1u_2) = 0$, then let $f'(u_2) := 5$, and if $f'(u_{2t}u_{2t+1}) \in \{5, 6\}$, then let $f'(u_{2t}) := 0$.

(4.2.3) Assume that $f'(u_1w_1) \neq 4$ and $f'(u_{2t+1}w_{2t+1}) = 1$.

Let $f' := f_8$.

First consider the case of $t = 3$. Let $f'(u_5u_7) := 0$. Next we assign a label in $\{5, 6\} - \{f'(u_1w_1)\}$ to (u_1, u_7) and a label in $\{0, 5, 6\} - \{f'(u_1w_1), f'(u_1u_7)\}$ to (u_1, u_2) . Reassign a label in $\{5, 6\} - \{f'(u_1u_7)\}$ to (u_6, u_7) . Then if $f'(u_1u_2) = 0$, then let $f'(u_2) := 5$.

Next consider the case of $t \geq 5$. Let $f'(u_{2t-2}) := 0$, $f'(u_{2t-1}) := 1$, $f'(u_{2t-2}u_{2t-1}) := 4$, $f'(u_{2t-1}u_{2t}) := 3$, and $f'(u_{2t-1}u_{2t+1}) := 5$. Next we assign a label in $\{0, 6\} - \{f'(u_1w_1)\}$ to (u_1, u_{2t+1}) and a label in $\{0, 5, 6\} - \{f'(u_1w_1), f'(u_1u_{2t+1})\}$ to (u_1, u_2) . Reassign a label in $\{0, 6\} - \{f'(u_1u_{2t+1})\}$ to (u_{2t}, u_{2t+1}) . Then if $f'(u_1u_2) = 0$, then let $f'(u_2) := 5$, and if $f'(u_{2t}u_{2t+1}) = 6$, then let $f'(u_{2t}) := 0$.

(4.2.4) Assume that $f'(u_1w_1) = 4$ and $f'(u_{2t+1}w_{2t+1}) = 1$.

Let $f' := f_8$. We reassign labels for some vertices and edges similarly to (4.2.2) and (4.2.3). Namely, if $t = 3$, let $f'(u_1u_2) := 0$, $f'(u_1u_3) := 6$, $f'(u_2) := 5$, $f'(u_6u_7) := 6$, $f'(u_5u_7) := 0$, and $f'(u_1u_7) := 5$. If $t \geq 5$, then let $f'(u_1u_2) := 5$, $f'(u_1u_3) := 6$, $f'(u_{2t-2}) := 0$, $f'(u_{2t-1}) := 1$, $f'(u_{2t}) := 0$, $f'(u_{2t-2}u_{2t-1}) := 4$, $f'(u_{2t-1}u_{2t}) := 3$, $f'(u_{2t}u_{2t+1}) := 6$, $f'(u_{2t-1}u_{2t+1}) := 5$, and $f'(u_1u_{2t+1}) := 0$. $\square$

References

1. F. Bazzaro, M. Montassier, and A. Raspaud. $(d,1)$ -total labelling of planar graphs with large girth and high maximum degree. *Discr. Math.* 307, 2141–2151 (2007).
2. M. Behzad. Graphs and their chromatic numbers. Ph.D. Thesis, Michigan State University, 1965.
3. T. Calamoneri. The $L(h,k)$ -labelling problem: A survey and annotated bibliography. *The Computer Journal* 49, 585–608 (2006). (The $L(h,k)$ -Labelling Problem: A Survey and Annotated Bibliography, <http://www.dsi.uniroma1.it/~calamo/PDF-FILES/survey.pdf>, ver. Jan. 13, 2009.)
4. D. Chen and W. Wang. $(2,1)$ -Total labelling of outerplanar graphs. *Discr. Appl. Math.* 155, 2585–2593 (2007).
5. S. Fiorini. On the chromatic index of outerplanar graphs. *J. Combin. Theory Ser. B* 18, 35–38 (1975).
6. J. R. Griggs and R. K. Yeh. Labelling graphs with a condition at distance 2. *SIAM J. Disc. Math.* 5, 586–595 (1992).
7. F. Havet and M. -L. Yu. $(d,1)$ -Total labelling of graphs. Technical Report 4650, INRIA, 2002.
8. F. Havet and M. -L. Yu. $(p,1)$ -Total labelling of graphs. *Discr. Math.* 308, 496–513 (2008).
9. D. König. Über graphen und ihre anwendug auf determinantentheorie und mengenlehre. *Math. Ann.* 77, 453–465 (1916).
10. K.-W. Lih, D. D.-F. Liu and W. Wang. On $(d, 1)$ -total numbers of graphs. *Discr. Math.* 309, 3767–3773 (2009).
11. M. Montassier and A. Raspaud. $(d,1)$ -total labeling of graphs with a given maximum average degree. *J. Graph Theory* 51, 93–109 (2006).
12. V. G. Vizing. On the estimate of the chromatic class of a p -graphs. *Diskret. Analiz.* 3, 25–30, (1964) (in Russian).
13. V. G. Vizing. Some unsolved problems in graph theory. *Russian Mathematical Surveys* 23, 125–141, (1968).
14. W. Wang and K. Zhang. Δ -Matchings and edge-face chromatic numbers. *Acta. Math. Appl. Sinica* 22, 236–242 (1999).
15. M. A. Whittlesey, J. P. Georges, and D. W. Mauro. On the λ -number of Q_n and related graphs. *SIAM J. Discr. Math.* 8, 499–506 (1995).
16. R. K. Yeh. A survey on labeling graphs with a condition at distance two. *Discr. Math.* 306, 1217–1231 (2006).